arXiv:1611.06411v1 [math.NT] 19 Nov 2016

Reductions of abelian varieties of generalized Mumford type

Steve Thakur

Abstract

We study the special fibers of a class of absolutely simple abelian varieties over number fields with endomorphism rings $\mathbb{Z}$ and l-adic monodromy groups of low rank. We also study the Dirichlet density of the places at which the possible reductions occur.

1 Introduction

In this article, we study abelian varieties of Mumford type and the Shimura curves parametrizing them, also known as generalized Mumford curves. First constructed in [Mum69], these are among the simplest examples of Shimura varieties that are of Hodge type but not of PEL type. Furthermore, they furnish us with examples of abelian varieties with trivial endomorphism rings such that the l-adic monodromy subgroups have the smallest possible rank. From [Orr15], it is known that for abelian varieties of dimension d and commutative endomorphism ring, we have the inequality

$$\text{rank } G_l(A) \geq \log_2 d + 2.$$

The abelian varieties we study in this article are examples where equality holds.

In particular, we study the isogeny classes of the reductions for dimension 16 (equivalently, rank 6), the dimension 4 (rank 4) case having been studied in [No00]. We also look into the densities of the primes where these reductions occur and confirm a special case of a broader conjecture for the splitting of reductions of abelian varieties with trivial endomorphism rings. Furthermore, for an abelian variety X of dimension g over a number field with $\text{End}(X) = \mathbb{Z}$, unless $2g$ lies in the (rather thin) set

$$\{n^{2k+1} : n, k \geq 1\} \bigcup \{\binom{4k+2}{2k+1} : k \geq 1\},$$

we have $G_l \cong \text{GSp}_{2g,\mathbb{Q}_l}$ and the Mumford-Tate conjecture holds for X [Pin98]. The simplest counter-examples are perhaps the ones we study in this article. Another reason we are interested in the dimension 16 case is that after [Zh14] that is the second smallest dimension (after 10) for which the Mumford-Tate conjecture remains open when the endomorphism ring is trivial (as far as we know).

Relying heavily on a combination of techniques from [Pin98], [No09] and [Zyw14], we deduce the following result:

Theorem 1.1. *Let A be an abelian variety of dimension 2^{2n} over a number field F such that:*
(1). The field F is enlarged so that the l-adic monodromy groups G_l are all connected
(2). For all primes l in a set of positive density, $\text{Lie}(G_l)_{\overline{\mathbb{Q}}_l} \cong \overline{\mathbb{Q}}_l \oplus \left(\bigoplus_{i=1}^{2n+1} \mathfrak{sl}_{2,\overline{\mathbb{Q}}_l}\right)$ with $\mathfrak{sl}_2^{2n+1}$ acting on $V_l \otimes_{\mathbb{Q}_l} \overline{\mathbb{Q}}_l$ by the $2n+1$-th power of the standard representation of $\mathfrak{sl}_2$.

Then the reduction A_v is absolutely simple away from a set of places of Dirichlet density zero.

This confirms a special case of the broader conjecture that absolutely simple abelian varieties over number fields with commutative endomorphism rings have absolutely simple reduction away from a set of places of density zero.

In the dimension 16 (equivalently, rank 6) case, we list the possible isogeny types for the reduction A_v (Proposition 4.8) and briefly describe how the same techniques may be used for higher dimensions when the rank of the derived subgroup of the l-adic monodromy group is a prime. In the case of good ordinary reduction - which occurs with density one - we show that the reduction is either simple or has an ordinary elliptic curve as a simple component. We also show that both types of reduction occur at some place for an abelian variety on the Mumford curve arising from a quaternion algebra over a totally real quintic number field.

1.1 Notations and Terminology

For an abelian variety A over a number field F and a prime l, the *l-adic monodromy subgroup* G_l is the Zariski closure of the image of $\rho_{_l} : \mathrm{Gal}_F \longrightarrow \mathrm{GL}(V_l(A))$ and G_l^0 is the connected component of the identity. We denote the Zariski closure of $\rho_l(\mathrm{Gal}_F)$ in $\mathrm{GL}(T_l(A))$ by $\mathcal{G}_l$. It is a group scheme over $\mathbb{Z}_l$ with generic fiber G_l.

Let g be the dimension of A. The Betti cohomology group $H^1(A(\mathbb{C}), \mathbb{Q})$ is a $2g$-dimensional vector space over $\mathbb{Q}$ endowed with a decomposition $V = V^{1,0} \oplus V^{0,1}$ such that $\overline{V^{1,0}} = V^{0,1}$. Let $\mu_\infty : \mathbb{G}_{m,\mathbb{C}} \longrightarrow \mathrm{GL}_{2g,\mathbb{C}}$ be the cocharacter through which any $z \in \mathbb{C}^*$ acts by multiplication by z on $V^{1,0}$ and trivially on $V^{0,1}$. The *Mumford-Tate group* $\mathrm{MT}(A)$ is the unique smallest $\mathbb{Q}$-algebraic subgroup of $\mathrm{GL}_{2g,\mathbb{Q}}$ such that μ_∞ factors through $\mathrm{MT}(A) \times_{\mathbb{Q}} \mathbb{C}$.

Conjecture (Mumford-Tate) For an abelian variety A over a number field and a prime l, $\mathrm{MT}(A) \times_{\mathbb{Q}} \mathbb{Q}_l = G_l^0$.

While the conjecture remains open, several cases have been proven. In particular, if the endomorphism ring is trivial, the two smallest dimensions for which the conjecture remains open are 10 and 16 ([Pin98], [Zh14]). Furthermore, the Mumford-Tate conjecture for abelian varieties over number fields is closely related to the simple decomposition of the special fibers of the abelian varieties ([Zyw14]).

The following is a straightforward generalization of the construction in [Mum69]. It is well-defined over an arbitrary field of characteristic zero. But throughout this article, we will only encounter cases where the field K is either a number field or a local field.

Definition 1.1. Let K be a field of characteristic zero, G an algebraic group over K and V a faithful K-linear representation of G. Then (G, V) is of *Mumford type of rank $N+1$* if:

-Lie(G) has a one-dimensional center
-$\mathrm{Lie}(G)_{\overline{K}} \cong \mathbb{G}_{a,\overline{K}} \oplus \underbrace{\mathfrak{sl}_{2,\overline{K}} \oplus \cdots \oplus \mathfrak{sl}_{2,\overline{K}}}_{N}$
-$\mathfrak{sl}_{2,\overline{K}}^N$ acts on $V_{\overline{K}}$ through the N-th tensor power of the standard representation of $\mathfrak{sl}_2$.

If G is also assumed to be connected, there is a central isogeny $N : \tilde{G} \longrightarrow G$ where $\tilde{G}$ is an algebraic group over K such that $\tilde{G} \cong \mathbb{G}_{m,K} \times \tilde{G}^{\mathrm{der}}$ and $\tilde{G}^{\mathrm{der}}_{\overline{K}} \cong \mathrm{SL}_{2,\overline{K}}^{2n+1}$

We introduce the two following notions so as to avoid ambiguity:

Definition 1.2. An abelian variety A over a number field F is *weakly* of Mumford type of rank N if (G_l, V_l) is of Mumford type of rank N for a set of rational primes l of density one.

Definition 1.3. An abelian variety A over a number field F is *strongly* of Mumford type of rank N if $(\mathrm{MT}(A), H^1_B(X(\mathbb{C}), \mathbb{Q}))$ is of Mumford type of rank N.

In both cases, the definitions imply that the abelian variety is of dimension 2^{2n}.

Let K be a totally real field, D a quaternion over K split at precisely one archimedean place with the corestriction $\mathrm{Cor}_{K/\mathbb{Q}}(D) = 0$ in $\mathrm{Br}(\mathbb{Q})$. Then an abelian variety corresponding to a closed point on the Shimura curve given by the datum $((D^*)^{\mathrm{der}}, h)$ fulfills the conditions of Definition 1.3. We refer the reader to ([SZ12], Section 2) for details. We will see in appendix 6.1 that the converse is also true.

An implication of ([Pin98], Theorem 5.15) is that an abelian variety *strongly* of Mumford type is *weakly* of Mumford type. The converse is known to be true for dimension 4 from [Zh14] and was the last remaining case of the Mumford-Tate conjecture for dimension 4. For higher dimensions, it remains an open case of the conjecture (as far as we know).

For an abelian variety A weakly of Mumford type, we fix a triple $(\mathbf{G}_A, \mathbf{V}_A, \mathbb{S}_A)$ such that:
- $\mathbf{G}_A$ is a connected quasi-split reductive group over $\mathbb{Q}$ with $\mathbf{G}_A^{\mathrm{der}}$ $\mathbb{Q}$-simple
- $\mathbf{V}_A$ is a faithful $\mathbb{Q}$-linear representation of $\mathbf{G}_A$
- $\mathbb{S}_A$ is a density one set of rational primes such that

$$(\mathbf{G}_A, \mathbf{V}_A) \otimes_{\mathbb{Q}} \mathbb{Q}_l \cong (G_l, V_l) \qquad \text{for all } l \in \mathbb{S}_A.$$

The existence of this triple follows from [Pin98, Theorem 5.13]. Furthermore, the pair $(\mathbf{G}_A, \mathbf{V}_A)$ is of Mumford type. If A is strongly of Mumford type, meaning that it arises as a closed point on a Mumford curve, the Mumford-Tate conjecture is true for A by [Pin98]. So the group $\mathbf{G}_A$ is, in fact, the Mumford-Tate group $\mathrm{MT}(A)$ and $\mathbf{V}_A$ is the Betti cohomology group $H^1_B(A(\mathbb{C}), \mathbb{Q})$.

2 Some background

In this section, we state some fundamental results that we shall need repeatedly.

Theorem 2.1 (Pin98)**.** *Let X be an abelian variety over a number field F such that $\mathrm{End}_{\overline{\mathbb{Q}}}(X) = \mathbb{Z}$ and the root system of each factor of $\mathrm{MT}(X)_{\overline{\mathbb{Q}}}$ has type A_{2s-1} with $s \geq 1$ or B_r with $r \geq 1$. Then the Mumford-Tate conjecture is true for X.*

In particular, if X is strongly of Mumford type, the root system of every factor is A_1 and the Mumford-Tate conjecture is true for X. So $G_l(X) = \mathrm{MT}(X) \times_{\mathbb{Q}} \mathbb{Q}_l$ and

$$\mathrm{Lie}(G_l(X))_{\overline{\mathbb{Q}}_l} \cong \mathrm{Lie}(\mathrm{MT}(X)) \otimes_{\mathbb{Q}} \mathbb{Q}_l \cong \mathbb{G}_{a,\overline{\mathbb{Q}}_l} \oplus \mathfrak{sl}^{2n+1}_{2,\overline{\mathbb{Q}}_l}.$$

Theorem 2.2. [LP95] *There exists a minimum finite extension F_A^{conn} where all the G_l are connected.*

This extension corresponds to the subgroup $\rho_l^{-1}(G_l^0(\mathbb{Q}_l)) \subseteq \mathrm{Gal}_F$ which is independent of l. We will usually be replacing F by F^{conn} in this article. This has the effect of replacing the groups G_l by their identity components. The field F_A^{conn} may be alternatively described as the intersection

$$F_A^{\mathrm{conn}} := \bigcap_l F(A[l^\infty]).$$

Theorem 2.3. (Faltings) *Let X be an abelian variety over a number field F such that $F_X^{\mathrm{conn}} = F$. Then we have the following:*
(i) *The centralizer of G_l in $\mathrm{End}_{\mathbb{Q}_l}(V_l)$ is $\mathrm{End}(X) \otimes_{\mathbb{Z}} \mathbb{Q}_l$.*
(ii) *The group G_l is reductive.*
(iii) $\mathrm{End}^0(X_F) = \mathrm{End}^0(X_{\overline{\mathbb{Q}}})$

The following result of Deligne gives one of the inclusions for the Mumford-Tate conjecture.

Theorem 2.4. (Deligne) *For any prime l, $G_l^0 \times_{\mathbb{Q}} \mathbb{Q}_l \subseteq \mathrm{MT}(A) \times_{\mathbb{Q}} \mathbb{Q}_l$.*

Theorem 2.5. (Serre) *For any abelian variety A, the l-adic monodromy groups $G_{l,A}$ are all of the same rank.*

Henceforth, we shall refer to this common rank as *the rank of the abelian variety.* Deligne's theorem on the first inclusion combined with Serre's rank independence theorem show that the Mumford-Tate conjecture reduces to the rank of the abelian variety being equal to the rank of the Mumford-Tate group. More precisely:

Theorem 2.6. (Serre) *If the inequality*

$$\text{rank } G_p^0 \leq \text{rank } \mathrm{MT}(A)$$

is an equality for any prime p, the Mumford-Tate conjecture is true for A.

Furthermore, if we impose the additional condition that the endomorphism ring is trivial, as is the case with abelian varieties weakly of Mumford type, the following theorem from [Pin98] shows there always exists a reductive group over $\mathbb{Q}$ that "looks like" the Mumford-Tate group is expected to.

Theorem 2.7. ([Pin98], Theorem 5.13) *Suppose* $\mathrm{End}(A) = \mathbb{Z}$. *Then there exists a quasi-split reductive group $\mathbf{G}_A \subseteq \mathrm{GL}_{2g,\mathbb{Q}}$ over $\mathbb{Q}$ with a faithful $\mathbb{Q}$-linear representation $\mathbf{V}_A$ such that*

$$(\mathbf{G}_A, \mathbf{V}_A) \otimes \mathbb{Q}_l \cong (G_l, V_l)$$

for all rational primes l in a set $\mathbb{S}_A$ of density one and $\mathbf{G}_A^{\mathrm{der}}$ is $\mathbb{Q}$-simple.

$\mathbf{G}_A^{\mathrm{der}}$ being $\mathbb{Q}$-simple implies that $\mathrm{Lie}(\mathbf{G}_A^{\mathrm{der}})$ is $\otimes$-isotypic. Furthermore, it is also known that the root systems of $\mathrm{Lie}(\mathbf{G}_A^{\mathrm{der}})$ are of type A, B, C or D ([Pin98]).

Theorem 2.8. ([LP95], Theorem 3.2) *For an abelian variety A over a field F such that $F = F_A^{\mathrm{conn}}$, G_l is unramified (quasi-split and split over some $\mathbb{Q}_{l^h}$) for all but finitely many l. Furthermore, the closed fiber of $\mathcal{G}_l$ is reductive over $\mathbb{Z}_l$ for all but finitely many l.*

(A reductive group over a field is *quasi-split* if it has a maximal torus that is quasi-split. If the field is a local field, the group is *unramified* if it is quasi-split and split over some unramified extension.)

2.1 Strict Compatibility and the Galois group of the characteristic polynomial

The l-adic representations attached to an abelian variety are *strictly compatible* in the sense of Serre. A proof may be found in [Del].

Theorem 2.9. [Del] *For an abelian variety A, fix a finite set $\mathcal{S}$ of non-archimedean places such that A has good reduction outside $\mathcal{S}$. Let $v \notin \mathcal{S}$, $l \neq \mathrm{char}(k_v)$*
(1) *ρ_l is unramified at v*
(2) *The characteristic polynomial of $\rho_l(\mathrm{Fr}_v)$ has coefficients in $\mathbb{Z}$ and is independent of l.*

An immediate consequence is that the eigenvalues are in $\overline{\mathbb{Z}}$ and are independent of l. We denote this polynomial by $P_{A_v}(X) \in \mathbb{Z}[X]$. Let $\pi_1, \cdots, \pi_{2g}$ be the eigenvalues of $\rho_l(\mathrm{Fr}_v)$. These are the zeros of $P_{A_v}(X)$. Let $\mathbb{Q}(\mathcal{W}_{A_v})^+$ be the compositum of all the totally real fields $\mathbb{Q}(\pi_i + \overline{\pi_i})$.

Then $\mathrm{Gal}(\mathbb{Q}(\mathcal{W}_{A_v})^+/\mathbb{Q}) \subseteq S_g$ and $\mathrm{Gal}(\mathbb{Q}(\mathcal{W}_{A_v})/\mathbb{Q}(\mathcal{W}_{A_v})^+)$ is abelian 2-torsion of rank at most g. Thus, the Galois group of $P_{A_v}(X)$ is a subgroup of the semi-direct product $W_{2g} := (\mathbb{Z}/2\mathbb{Z})^g \rtimes S_g$, the "generic Galois group" of the Galois closure of a CM field of degree $2g$.

The Galois group is the full W_{2g} away from a density zero set of places when the endomorphism ring is trivial and $G_l = \mathrm{GSp}_{\mathbb{Q}_l}$ [Bog80]. On the other hand, the Galois group is far smaller for abelian varieties weakly of Mumford type on account of the additive relations between the weights of the representation of $G_{l,\overline{\mathbb{Q}_l}}$ which induce multiplicative relations between the eigenvalues of the Frobenius.

3 Some preliminary results

In this section, we discuss a few results we shall need in the subsequent sections. Some of them will be necessary to verify that the concepts from the preceding section are well-defined.

Proposition 3.1. *Let A be an abelian variety of dimension 2^{2n} over a number field F. Suppose there exists a prime p in $\mathbb{S}_A$ such that (G_p, V_p) is of Mumford type. Then* $\mathrm{End}(A_{\overline{\mathbb{Q}}}) = \mathbb{Z}$

Proof. Passing to a finite extension if necessary, we may assume without loss of generality that all the l-adic monodromy groups are connected. Now, by Faltings,

$$\mathrm{End}^0(A_{F'}) \otimes_{\mathbb{Q}} \mathbb{Q}_p \cong \mathrm{End}_{\mathrm{Gal}_{F'}}(V_p) = \mathrm{End}_{G_p}(V_p) = \mathbb{Q}_p$$

for any finite extension F'/F. Thus, $\mathrm{End}^0(A_{\overline{\mathbb{Q}}}) = \mathbb{Q}$. □

Proposition 3.2. *Let A be an abelian variety of dimension 2^{2n} over a number field. Suppose the set of rational primes p such (G_p, V_p) is of Mumford type of rank $2n+2$ has positive density. Then (G_l, V_l) is of Mumford type for all $l \in \mathbb{S}_A$.*

Proof. By [Pin98], there exists a reductive quasi-split group $\mathbf{G}_A$ over $\mathbb{Q}$ with a faithful $\mathbb{Q}$-linear representation $\mathbf{V}_A$ such that $(\mathbf{G}_A, \mathbf{V}_A) \otimes_{\mathbb{Q}} \mathbb{Q}_l \cong (G_l, V_l)$ for all $l \in \mathbb{S}_A$. Since the set of rational primes described in the Proposition has positive density, it intersects $\mathbb{S}_A$. So there exists a prime p such that $(\mathbf{G}_A, \mathbf{V}_A) \otimes \mathbb{Q}_p \cong (G_p, V_p)$ and (G_p, V_p) is of Mumford type. Hence, $(\mathbf{G}_A, \mathbf{V}_A)$ is of Mumford type of rank $2n+2$ and the result follows. □

In the dimension 4 case, (G_p, V_p) being of Mumford type for a single prime p implies the same for all primes. We refer the reader to [No00] for details. The proof follows from Serre's rank independence result and the classification of semisimple Lie algebras.

Lemma 3.3. *Let G be a $\mathbb{Q}$-simple algebraic group. If G is semisimple and the number of simple factors of $G_{\overline{\mathbb{Q}}}$ is a prime p, then the set of primes l such that $G_{\mathbb{Q}_l}$ is $\mathbb{Q}_l$-simple is of positive density (in particular, infinite).*

Proof. The action of $\mathrm{Gal}_{\mathbb{Q}}$ on the simple factors of $G_{\overline{\mathbb{Q}}}$ gives a map $\tau : \mathrm{Gal}_{\mathbb{Q}} \to S_p$. Since G is $\mathbb{Q}$-simple, the image of τ is a transitive subgroup of S_p and hence, contains a p-cycle c. By the Chebotarev density theorem, the set of primes l such that $c \in \tau(\mathrm{Gal}_{\mathbb{Q}_l})$ has positive density and in particular, is infinite. For any $l \in \mathbb{S} \cap \mathcal{P}$, $\mathrm{Gal}_{\mathbb{Q}_l}$ acts transitively on the simple factors of $G_{\overline{\mathbb{Q}}_l}$ and hence, $G_{\mathbb{Q}_l}$ is $\mathbb{Q}_l$-simple. □

Lemma 3.4. *Let A be an abelian variety over number field such that* $\mathrm{End}(A_{\overline{\mathbb{Q}}}) = \mathbb{Z}$. *Suppose the number of simple factors of $G^{\mathrm{der}}_{l,\overline{\mathbb{Q}}}$ is a prime p for all l in a set of density one. Then there exist infinitely many primes l such that G^{der}_l is $\mathbb{Q}_l$-simple.*

Proof. From [Pin98], there exists a connected reductive group $\mathbf{G}_A$ over $\mathbb{Q}$ with a faithful linear representation $\mathbf{V}_A$ such that:

-$(\mathbf{G}_A, \mathbf{V}_A) \otimes \mathbb{Q}_l \cong (G_l, V_l)$ for all l in a set $\mathbb{S}_A$ of density one
-$\mathbf{G}_A^{\mathrm{der}}$ is $\mathbb{Q}$-simple

Now $\mathrm{Lie}(\mathbf{G}_A)$ has a one-dimensional center and $\mathrm{Lie}(\mathbf{G}_A^{\mathrm{der}})$ is $\mathbb{Q}$-simple with $\mathrm{Lie}(\mathbf{G}_{A,\overline{\mathbb{Q}}}^{\mathrm{der}})$ having p simple factors. The permutation action of $\mathrm{Gal}_{\mathbb{Q}}$ on the simple factors of $\mathbf{G}_{A,\overline{\mathbb{Q}}}^{\mathrm{der}}$ gives a map $\tau : \mathrm{Gal}_{\mathbb{Q}} \to S_p$. Since $\mathbf{G}_A^{\mathrm{der}}$ is $\mathbb{Q}$-simple, it follows that the image is a transitive subgroup and since p is a prime by assumption, the image contains a p-cycle c. But $c \in \tau(\mathrm{Gal}_{\mathbb{Q}_l})$ for a set $\mathcal{P}$ of rational primes of positive density by the Chebotarev density theorem. Since $\mathbb{S}_A$ is of density one, the intersection $\mathbb{S}_A \cap \mathcal{P}$ is also of positive density. For any prime $l \in \mathbb{S}_A \cap \mathcal{P}$, G_l^{der} is $\mathbb{Q}_l$-simple since the p-cycle acts transitively on the simple factors of $G_{l,\overline{\mathbb{Q}}}^{\mathrm{der}}$. □

We now use this lemma to show potential good reduction in the case when $2n+1$ is prime. The key idea (following [No00]) is to use the fact that any abelian variety is De Rham everywhere, meaning that the inertia group acts unipotently on the Tate module of every different prime after passing to a finite extension, and subsequently show that the inertia group acts trivially on the Tate module.

Proposition 3.5. *Suppose $2n+1$ is a prime. Let A be an abelian variety of rank $2n+2$ weakly of Mumford type over a number field F. Then A has potential good reduction everywhere.*

Proof. Enlarging the field if necessary, we may assume without loss of generality that all the G_l are connected and A has semistable reduction everywhere. Since $2n+1$ is assumed to be prime, there exist infinitely many primes l such that G_l^{der} is $\mathbb{Q}_l$-simple.

Let v be a finite place of F and let I_v be the inertia group. Let p be the rational prime lying under v. Choose a prime $l \neq p$ such that G_l^{der} is $\mathbb{Q}_l$-simple and let ρ_l be the l-adic representation associated to A. Since A is semistable at v, the inertia group I_v is rank 2 unipotent on $V_l(A)$, meaning that $(\rho_l(\sigma) - \mathrm{id})^2 = 0$ for every $\sigma \in I_v$ ([GRR72]).

Suppose, by way of contradiction, that $\rho_l(\sigma) \neq \mathrm{id}$ for some $\sigma \in I_v$. Now $\rho_l(\sigma) \in G_l(\overline{\mathbb{Q}})$ is the image by tensor product of an element $(y_1, \cdots, y_{2n+1}) \in \mathrm{SL}_2(\overline{\mathbb{Q}}_l)$ with at least one $y_i \neq \mathrm{id}$. But since G_l^{der} is $\mathbb{Q}_l$-simple, it follows that $y_i \neq \mathrm{id}$ for any i. Conjugating if necessary, we may assume without loss of generality that all the y_i are of the shape

$$\begin{pmatrix} 1 & a_i \\ 0 & 1 \end{pmatrix}$$

with each $a_i \in \overline{\mathbb{Q}}_l \setminus \{0\}$. So the index of nilpotency of $x - \mathrm{id}$ is larger than 2, a contradiction. Hence, I_v acts trivially on V_l after passing to a finite extension. From the Neron-Ogg-Shafarevich criterion ([ST68]), it follows that A has good reduction at v. □

For abelian varieties *strongly* of Mumford type, [SZ12] have a proof of potential good reduction that does not assume $2n+1$ being a prime. They do, however, assume that v is prime to the discriminant of the totally real field of degree $2n+1$ and to the discriminant of the quaternion algebra over that field that yields the Shimura curve.

Furthermore, ([Pin98], Theorem 7.1) implies that the reduction is ordinary away from a density zero set.

Proposition 3.6. *For an abelian variety A weakly of Mumford type over a field $F = F_A^{\mathrm{conn}}$, the places of good ordinary reduction have Dirichlet density* 1.

Proof. The root system of $G^{\mathrm{der}}_{l,\overline{\mathbb{Q}}_l}$ has type A_1 for a set of primes of density one. So the same is true for the root system of $\mathbf{G}^{\mathrm{der}}_{A,\overline{\mathbb{Q}}}$. In particular, the simple factors of $\mathbf{G}^{\mathrm{der}}_{A,\overline{\mathbb{Q}}}$ are not of type C_r for any $r \geq 3$ and hence, A has good ordinary reduction away from a density zero set of places by ([Pin98], Theorem 7.1). □

An alternate proof independent of the results of [Pin98] may be found in [No95].

Proposition 3.7. *If an abelian variety X of dimension 2^{2n} is strongly of Mumford type, then it is weakly of Mumford type.*

Proof. By the hypothesis, the roots system of every simple factor of $\mathrm{MT}(X)_{\overline{\mathbb{Q}}}$ has type A_1. So, by ([Pin98], Theorem 5.15), the Mumford-Tate conjecture is true for X. Hence,

$$\mathrm{Lie}(G_{X,l})_{\overline{\mathbb{Q}}_l} \cong \mathrm{Lie}(\mathrm{MT}(X) \otimes_{\mathbb{Q}} \mathbb{Q}_l)_{\overline{\mathbb{Q}}_l} = \overline{\mathbb{Q}}_l \oplus \mathfrak{sl}^{2n+1}_{2,\overline{\mathbb{Q}}_l}.$$

Alternatively, note that if X is strongly of Mumford type, then rank $\mathrm{MT}(X) = 2n + 2$. By [Orr15], we have the inequality rank $G_{X,l} \geq 2n+2$. Hence, rank $G_{X,l} \geq$ rank $\mathrm{MT}(X)$, which by Theorem 2.6, implies the Mumford-Tate conjecture for X. □

For the next two lemmas, let A be an absolutely simple abelian variety over a number field F. Let v be a place such that $\mathrm{End}(A_{\overline{k}})$ is commutative. Note that the set of such places has density one if $\mathrm{End}(A_{\overline{\mathbb{Q}}})$ is commutative ([LP97]) and is empty if $\mathrm{End}(A)$ is non-commutative.

Lemma 3.8. *With this setup, none of the quotients $\frac{\pi_1}{\pi_2}$ of the eigenvalues of Fr_v are roots of unity.*

Proof. Suppose Fr_v has two eigenvalues π_1, π_2 such that $\frac{\pi_1}{\pi_2} = \zeta_N$ for some integer N. Choose a finite extension F'/F and a place $\tilde{v}$ lying over v with inertia degree N over v. Then $\mathcal{W}_{A_{\tilde{v}}} = \mathcal{W}^N_{A_v}$ and hence, $P_{A_{\tilde{v}}}(X)$ has the zero π_1^N with multiplicity at least two. In particular, $P_{A_{\tilde{v}}}(X)$ is inseparable and hence, $\mathrm{End}(A_{\tilde{v}})$ is non-commutative, a contradiction. □

Lemma 3.9. *The reduction A_v is absolutely simple if and only if $P_{A_v}(X)$ is irreducible.*

Proof. Since $\mathrm{End}(A_{\overline{k}})$ is commutative, the zeros of $P^N_{A_v}(X)$ are distinct. So A_v is absolutely simple if and only $\mathrm{Gal}_{\mathbb{Q}}$ acts transitively on $\mathcal{W}^N_{A_v} = \mathcal{W}_{A_v \otimes k_{v^N}}$ for every integer N. Let $\sigma \in \mathrm{Gal}_{\mathbb{Q}}$ and let π_1, π_2 be any two eigenvalues of Fr_v. Then $\sigma(\pi_1^N) = \pi_2^N \iff \left(\frac{\sigma(\pi_1)}{\pi_2}\right)^N = 1 \iff \sigma(\pi_1) = \pi_2$ by the preceding lemma. So $\mathrm{Gal}_{\mathbb{Q}}$ acts transitively on $\mathcal{W}^N_{A_v}$ if and only if it acts transitively on $\mathcal{W}_{A_v}$, which is equivalent to $P_{A_v}(X)$ being irreducible. □

The lemmas 3.8 and 3.9 will help us detect places of absolutely simple reduction. In particular, as we will see in section 4, for an abelian variety weakly of Mumford type of rank $2n+2$ with the integer $2n+1$ a prime, absolutely simple reduction only occurs when $\mathrm{End}(A_{\overline{k}})$ is commutative.

4 Isogeny Classes

We now study the isogeny classes of the special fibers of abelian varieties of dimension 16 (equivalently, rank 6) weakly of Mumford type. In particular, we determine the possible dimensions

of the simple factors of the reductions and their endomorphism algebras. We will follow some of the techniques from [No00] but will need a few more techniques than in the dimension 4 case. We also briefly describe how the techniques could be adapted to certain higher dimensional cases.

4.1 The Galois group of the characteristic polynomial

Some Notation: Let A be an abelian variety of rank $2n+2$ weakly of Mumford type over a number field F. Let v be a place of good reduction and let l be a prime other than the residue characteristic of v such that (G_l, V_l) is of Mumford type. Let $\rho_l : \mathrm{Gal}_F \longrightarrow \mathrm{GL}_{2g}(\mathbb{Z}_l)$ be the l-adic Galois representation and Fr_v a geometric Frobenius. Let $\pi_1, \cdots, \pi_{2g}$ be the eigenvalues of $\rho_l(\mathrm{Fr}_v)$. The eigenvalues lie in $\overline{\mathbb{Z}}$ and are independent of l. We will denote the set of eigenvalues by $\mathcal{W}_{A_v}$ and the multiplicative group they generate by $\Phi_{A_v} \subseteq \overline{\mathbb{Q}}^*$. The irreducible decomposition of the polynomial $P_{A_v}(X) = \prod_{i=1}^{2g}(X - \pi_i) \in \mathbb{Z}[X]$ determines the simple decomposition of A_v. In particular, $P_{A_v}(X)$ is separable if and only if $\mathrm{End}^0(A_v)$ is a CM algebra, as is the case when v is a place of ordinary reduction. Furthermore, $P_{A_v}(X)$ is irreducible if and only if A_v is simple. The field $\mathbb{Q}(\mathcal{W}_{A_v})$ is a Galois extension of $\mathbb{Q}$ and

$$\mathrm{Gal}(\mathbb{Q}(\mathcal{W}_{A_v})/\mathbb{Q}) \subseteq \mathrm{Aut}\,(X^*(\mathbf{T})) = \{\pm 1\}^{2n+1} \rtimes S_{2n+1}$$

where $\mathbf{T}$ is a maximal torus of $\mathbf{G}_A$. If F'/F is a finite extension and v' is a place of F' lying over v, we have $\Phi_{A_{v'}} = \Phi_{A_v}^d$ where d is the inertia degree of v' over v.

If the integer $2n+1$ is a prime, $\mathrm{Gal}(\mathbb{Q}(\mathcal{W}_{A_v})/\mathbb{Q}) = \{\pm 1\}^j \rtimes H$ with H a subgroup of S_{2n+1} containing a $2n+1$-cycle and $1 \leq j \leq 2n+1$. Also, it is easy to show that $2^j \equiv 1 \pmod{2n+1}$ (Proposition 4.10).

4.2 Weil Numbers and Isokummerian abelian varieties

We first state a few long-known results that will be useful for determining the possible isogeny classes of the simple components of the reductions. Proofs may be found in [Tat69] and [Wat].

Notation: For a Weil q-integer π (upto conjugacy), let B_π be the corresponding simple abelian variety over $\mathbb{F}_q$ (upto isogeny).

Lemma 4.1. *Let π be a Weil q-integer and B_π be the corresponding abelian variety over $\mathbb{F}_q$. The following are equivalent:*
(1) *B_π is supersingular, meaning that the Newton slopes are all $\frac{1}{2}$*
(2) *$\pi^N \in \mathbb{Q}$ for some integer N*
(3) *$\mathrm{End}^0_{\mathbb{F}_p}(B_\pi) = \mathrm{End}^0_{\mathbb{F}_{q^2}}(B_\pi) \cong \mathrm{Mat}_g(\mathbb{Q}_{p,\infty})$ where $\mathbb{Q}_{p,\infty}$ is the quaternion over $\mathbb{Q}$ ramified only at p and ∞.*

We may enlarge the field so that the size q of the residue field is a square and none of the quotients $\frac{\pi}{\sqrt{q}}$ are non-trivial roots of unity. So the only possible Weil q-integer corresponding to a supersingular abelian variety would be $\sqrt{q}$.

Lemma 4.2. *For a simple abelian variety B_π over $\mathbb{F}_q$, the following are equivalent:*
1. *$\mathrm{End}^0(B_\pi)$ is commutative.*
2. *The Weil number π is of degree $2\dim B_\pi$.*

Lemma 4.3. *For a simple abelian variety B_π over $\mathbb{F}_q$, the following are equivalent:*
1. *B_π is ordinary, meaning that the p-torsion group scheme is of rank g.*

2. $(\pi, \overline{\pi}) = (1)$ *in the ring of integers of the Galois closure of* $\mathbb{Q}(\pi)$.
2.′ $(\pi + \overline{\pi}, q) = (1)$ *in the ring of integers of the Galois closure of* $\mathbb{Q}(\pi)$.
3. *The Newton polygon is* $g \times 0, g \times 1$

The group Φ_{A_v}. For an abelian variety B over a finite field, we denote the multiplicative group generated by the eigenvalues of the Frobenius by Φ_B. For a Weil number π, Φ_π is the group Φ_{A_π} where A_π is an abelian variety in the corresponding isogeny class. So Φ_π is the $\mathrm{Gal}_{\mathbb{Q}}$-stable subgroup of $\overline{\mathbb{Q}}^*$ generated by the conjugates of π over $\mathbb{Q}$.

We introduce the following notion that is weaker than isogeny between two abelian varieties over finite fields.

Definition 4.1. Following [Kow03], we say two abelian varieties B_1, B_2 over a finite field are *isokummerian* if there exists some integer N such that $k(B_1[d]) = k(B_2[d])$ for all d prime to N.

The following lemma establishes an equivalence between this notion and that of the the multiplicative groups generated by the eigenvalues coinciding.

Proposition 4.4. ([Kow03], Lemma 8.1) *Two abelian varieties B_1, B_2 over a finite field are isokummerian if and only if $\Phi_{B_1} = \Phi_{B_2}$.*

Clearly, two isogenous abelian varieties are isokummerian. The converse is true, for instance, when one of the two is an elliptic curve (on account of the lack of non-torsion units in imaginary quadratic fields). But it is not true in general even for ordinary abelian varieties, as evidenced by a class of counterexamples constructed in ([Kow06], Remark 2.5). However, for Weil numbers associated to ordinary abelian varieties, we have the following result.

Proposition 4.5. ([Kow06], Proposition 2.1) *Let B be an abelian variety of dimension ≥ 3 over a finite field k of size q. Suppose the corresponding Weil number π is such that* $\gcd(\pi, \overline{\pi}) = (1)$ *in the Galois closure of* $\mathbb{Q}(\pi)$. *Suppose, further, that the exponent 2 abelian Galois group* $\mathrm{Gal}(\mathbb{Q}(\mathcal{W}_B)/\mathbb{Q}(\mathcal{W}_B)^+)$ *is of rank* $\dim B$. *Then the only Weil q-integers in the multiplicative group Φ_B are the conjugates of π.*

The next two lemmas will be useful for detecting places of ordinary reduction. As mentioned earlier, if B_π is ordinary, the endomorphism algebra is commutative ([Tat69]). While the converse is certainly false, we have the following:

Lemma 4.6. *Let B_π be an absolutely simple abelian variety over a finite field k. Suppose:*
(i) $\mathrm{End}^0_k(B_\pi)$ *is commutative*
(ii) p splits completely in $\mathbb{Q}(\pi)$.
Then B_π is ordinary.

Proof. The Newton slopes of B_π are given by $\frac{w(\pi)}{w(q)}$, where w runs through all the places of $\mathbb{Q}(\pi)$ lying over p. Since $\mathrm{End}^0(B_\pi)$ is commutative, it follows from Honda-Tate theory that the local invariants $[\mathbb{Q}(\pi)_w : \mathbb{Q}_p]\frac{w(\pi)}{w(q)}$ are integers. But since p splits completely in $\mathbb{Q}(\pi)/\mathbb{Q}$ and hence, in the Galois closure $\mathbb{Q}(\mathcal{W}_{B_\pi})$, any prime of $\mathbb{Q}(\mathcal{W}_{B_\pi})$ lying over p has local degree 1 over $\mathbb{Q}$. So the Newton slopes $\frac{w(\pi)}{w(q)}$ are integers and hence, B_π is ordinary. □

Lemma 4.7. *Let B_π be an absolutely simple abelian variety over a finite field k. Suppose:*
(i) $\mathrm{End}^0_k(B_\pi)$ *is commutative*
(ii) The Galois group $\mathrm{Gal}(\mathbb{Q}(\mathcal{W}_{B_\pi})/\mathbb{Q})$ *is such that the decomposition group of a prime of* $\mathbb{Q}(\mathcal{W}_{A_v})$ *lying over p is a normal subgroup of* $\mathrm{Gal}(\mathbb{Q}(\mathcal{W}_{B_\pi})/\mathbb{Q})$.
Then B_π is ordinary.

Remark In particular, the second condition is fulfilled when the center $\mathbb{Q}(\pi)$ of $\mathrm{End}^0(B_\pi)$ is an abelian extension over $\mathbb{Q}$.

Proof. Let $\mathfrak{p}$ be a prime of $\mathbb{Q}(\mathcal{W}_{B_\pi})$ lying over p. If $\mathfrak{p}$ has local degree 1, then clearly all the slopes are integral and hence, must be either 0 or 1. So we may assume without loss of generality that p does not split completely in $\mathbb{Q}(\mathcal{W}_{B_\pi})$.

Choose a non-trivial element τ in the decomposition group $D_\mathfrak{p}$. Since π and its conjugates generate $\mathbb{Q}(\mathcal{W}_{B_\pi})$, τ cannot fix every conjugate of π. So we may assume without loss of generality that $\tau(\pi) \neq \pi$. Furthermore, the hypothesis implies that $D_\mathfrak{p}$ is normal. So τ lies in the decomposition group of every prime of $\mathbb{Q}(\mathcal{W}_{A_v})$ lying over p and hence, τ preserves the ideal (π) in $\mathbb{Q}(\mathcal{W}_{B_\pi})$. So the algebraic number $\frac{\tau(\pi)}{\pi}$ has the valuation 0 under every non-archimedean valuation and hence, must be a root of unity. Write $\tau(\pi) = \zeta_N \pi$. Then the endomorphism algebra $\mathrm{End}^0(B \times_{\mathbb{F}_q} \mathbb{F}_{q^N})$ is non-commutative since the characteristic polynomial of $B \times_{\mathbb{F}_q} \mathbb{F}_{q^N}$ has the zero π^N with multiplicity ≥ 2, a contradiction. So p splits completely in $\mathbb{Q}(\mathcal{W}_{B_\pi})/\mathbb{Q}$ and by the preceding lemma, it follows that B_π is ordinary. □

4.3 Newton Polygons

With techniques virtually identical to those of ([No00], Section 3), we can show that for an abelian variety weakly of Mumford type of rank 6, the Newton slopes are given by

$$\frac{1}{2} \pm a_1 \pm a_2 \pm a_3 \pm a_4 \pm a_5$$

where the a_i are rational numbers between 0 and 1. Using the fact that the Newton slopes lie between 0 and 1, have integral break points and the Newton polygons of abelian varieties are admissible, we can then deduce that the slope $\frac{1}{2}$ occurs with multiplicity $0, 8, 12$ or 16. This will be be useful in the proof of Proposition 4.8.

4.4 Reductions of abelian varieties weakly of Mumford type

Let A be a 16-dimensional (equivalently, rank 6) abelian variety weakly of Mumford type over a number field F. Enlarging F if necessary, we may assume without loss of generality that A has good reduction everywhere. Let v be a non-archimedean place with $p = \mathrm{char}(v)$. We choose a prime $l \neq p$ such that the derived subgroup G_l^{der} is $\mathbb{Q}_l$-simple and (G_l, V_l) is of Mumford type. As seen in lemma 3.4, the set of primes fulfilling the first condition has positive density and since $\mathbb{S}_A$ has density one, the intersection has positive density as well. In particular, there are infinitely such primes.

Proposition 4.8. *With the setup as above, the reduction $A_{\overline{k}}$ has one of the following simple decompositions upto isogeny:*

(1) $(A^{(1)})^{16}$ *where $A^{(1)}$ is a supersingular elliptic curve*
(2) $A^{(1)} \times A_1^{(5)} \times A_2^{(5)} \times A_2^{(5)}$ *with Newton polygon* $0, 5 \times \frac{1}{5}, 10 \times \frac{2}{5}, 10 \times \frac{3}{5}, 5 \times \frac{4}{5}, 1$
(3) $A^{(1)} \times A^{(5)} \times A^{(10)}$ *with $A^{(1)}$ supersingular*
(4) $A^{(1)} \times A^{(5)} \times A^{(10)}$ *with $A^{(1)}$ ordinary*
(5) $A^{(1)} \times A_1^{(5)} \times A_2^{(5)} \times A_3^{(5)}$ *with each component ordinary and the last two isokummerian*
(6) *Simple*

where the superscript is the dimension of the simple abelian variety and distinct subscripts indicate non-isogeny. Furthermore, in the last three cases, $\mathrm{End}(A_{\overline{k}})$ *is commutative. The last possibility occurs with Dirichlet density one.*

Remark In the course of this proof, we will also determine the possibilities for the endomorphism algebra $\mathrm{End}^0(A_{\overline{k}})$. That is a more concrete description of the simple decomposition since the dimension of a simple abelian variety can be easily retrieved from the endomorphism algebra via Honda-Tate theory. Furthermore, we also look into the Galois group $\mathrm{Gal}(\mathbb{Q}(\mathcal{W}_{A_v})/\mathbb{Q})$ which provides some insight into the reduction A_v.

Proof. Let

$$\rho_{v,l} : \mathrm{Gal}_{F_v} \cong D_v \hookrightarrow \mathrm{Gal}_F \longrightarrow G_l(\mathbb{Z}_l)$$

be the local l-adic Galois representation attached to A. Replacing F by a finite extension if necessary, we may assume the following:

(1) $q := |k|$ is a perfect square.
(2) All the l-adic monodromy groups are connected.
(3) All elements of $\rho_{v,l}(\mathrm{Gal}_{F_v})$ are congruent to the identity mod l^2.
(4) All simple factors of $A_{\overline{k}}$ are defined over k.

The first and fourth condition make the Honda-Tate classification easier to work with. In particular, the simple components of A_v are, in fact, absolutely simple. So for any Weil number π corresponding to a simple component of A_v, the inclusion $\mathbb{Q}(\pi^d) \subseteq \mathbb{Q}(\pi)$ is an equality for any integer d. The second implies that any element of $\rho_{v,l}(\mathrm{Gal}_{F_v})$ lifts uniquely to an element of $\tilde{G}$. The third ensures that the group generated by the eigenvalues of the Frobenius does not contain any roots of unity other than 1.

Now, there exists a universal simple cover $N : \tilde{G}_l \longrightarrow G_l$ with a central isogeny N. From the third assumption above, $\rho_{v,l}$ lifts uniquely to a representation $\tilde{\rho}_{v,l} : \mathrm{Gal}_{F_v} \longrightarrow \tilde{G}_l(\mathbb{Z}_l)$ with the same congruence condition. Because of the uniqueness of the lift, $\tilde{\rho}_{v,l}$ is a group homomorphism.

Let $\mathrm{Fr_v}$ be a geometric Frobenius. Set $\pi = \tilde{\rho}_{v,l}(\mathrm{Fr_v}) \in \tilde{G}_l(\mathbb{Q}_l)$ and define $\widetilde{T} \subseteq \tilde{G}_l$ as the Zariski closure of the subgroup of $\tilde{G}_l(\mathbb{Q}_l)$ generated by π. The congruence condition on $\tilde{\rho}$ ensures that the subgroup of $\overline{\mathbb{Q}}^*$ generated by the eigenvalues of π does not contain any roots of unity other than 1 and hence, $\widetilde{T}$ is connected. So $\widetilde{T}$ is a maximal torus in $\tilde{G}_l$.

Now $\sqrt{q} \in \mathbb{Z}$ by construction. Set $\alpha := \frac{\pi}{\sqrt{q}} \in \tilde{G}_l(\mathbb{Q}_l)$ and let $\widetilde{T}'$ be the Zariski closure of the subgroup of $\tilde{G}_l$ generated by α. Then $\widetilde{T} \cong \mathbb{G}_{m,\mathbb{Q}_l} \times \widetilde{T}'$ with $\widetilde{T}'$ a maximal torus in $\tilde{G}^{\mathrm{der}}$. We choose a maximal torus $\overline{T}_{\mathbb{Q}_l}$ of ${\tilde{G}_l}^{\mathrm{der}}$ (over $\mathbb{Q}_l$) containing $\widetilde{T}'$. Since (G_l, V_l) is of Mumford type of rank 6, $\overline{T}$ is of rank 5.

Let Ψ be the associated root system of $\widetilde{T}'$. Let $\mathrm{Out}(\Psi) = \mathrm{Aut}(\Psi)/W(\Psi)$ be the group of outer automorphism group of Ψ, which is isomorphic to S_5. There exists an isomorphism $X^*(\overline{T}) \cong \mathbb{Z}^5$ such that the weights of the representation of $\overline{T}$ on V_l correspond to the elements $(\pm 1, \cdots, \pm 1) \in \mathbb{Z}^5$. The action of $\mathrm{Gal}_{\mathbb{Q}_l}$ on $X^*(\overline{T})$ induces an action on $\mathbb{Z}^5$. This action factors through the group $\{\pm 1\}^5 \rtimes S_5$ and contains a 5-cycle, which we may assume without loss of generality to be the cycle (1 2 3 4 5).

Evaluation at the element $\alpha \in \overline{T}(\mathbb{Q}_l)$ defines a $\mathrm{Gal}_{\mathbb{Q}_l}$-equivariant map

$$\mathrm{ev} : X^*(\overline{T}) \longrightarrow \overline{\mathbb{Q}}_l^*, \quad \chi \mapsto \chi(\alpha).$$

But the characteristic polynomial of the Frobenius lies in $\mathbb{Z}[X]$ and is independent of l. Hence, the eigenvalues of $\rho_{v,l}(\mathrm{Fr}_v)$ lie in $\overline{\mathbb{Q}}$ where $\overline{\mathbb{Q}}$ is some algebraic closure of $\mathbb{Q}$ embedded in $\overline{\mathbb{Q}}_l$. Since the weights of the representation of $\overline{T}$ on V_l correspond to the elements $(\pm 1, \pm 1, \pm 1, \pm 1, \pm 1) \in \mathbb{Z}^5$, the images $\mathrm{ev}(\pm 1, \cdots, \pm 1)$ are the eigenvalues of α on V_l. It follows that $\mathrm{ev}(\pm 1, \cdots, \pm 1)$ are

in $\overline{\mathbb{Q}} \subseteq \overline{\mathbb{Q}}_l$. So the image of ev lies in $\overline{\mathbb{Q}}^*$. As $\widetilde{T}'$ is defined as the Zariski closure of the subgroup of $\tilde{G}_l(\mathbb{Q}_l)$ generated by α, it follows that $\ker(\mathrm{ev})$ is the kernel of the natural surjection $X^*(\overline{T}) \longrightarrow X^*(\widetilde{T}')$. So we may identify $X^*(\widetilde{T}')$ with $\mathbb{Z}^5/\ker(\mathrm{ev})$.

For any element $(x_1, \cdots, x_5)$, denote its image in $X^*(\widetilde{T}')$ by $(x_1, \cdots, x_5)'$. The map ev induces an action of $\mathrm{Gal}_{\mathbb{Q}}$ on $X^*(\widetilde{T}')$ extending the action of $\mathrm{Gal}_{\mathbb{Q}_l}$. This action stabilizes the elements $\{(\pm 1, \cdots, \pm 1)\} \subseteq X^*(\widetilde{T}')$. Since all the eigenvalues of α have absolute value 1, it follows that $\mathrm{ev}(P)\overline{\mathrm{ev}(P)} = 1$ for each vertex P of the hypercube $(\{\pm 1, \cdots, \pm 1\})$. So the complex conjugation acts by multiplication by -1 on $X(\widetilde{T}')$.

We fix an embedding $\overline{\mathbb{Q}} \hookrightarrow \overline{\mathbb{Q}}_p$ which induces a p-adic valuation $\mathfrak{v}$ on $\overline{\mathbb{Q}}$ normalized by $\mathfrak{v}(q) = 1$ and define $\phi = \mathfrak{v} \circ \mathrm{ev}' : X^*(\widetilde{T}') \longrightarrow \mathbb{Q}$. This map is $\mathbb{Z}$-linear but not necessarily $\mathrm{Gal}_{\mathbb{Q}}$-equivariant or injective. The Newton slopes are then given by $\frac{1}{2} + \phi(\pm 1, \pm 1, \pm 1, \pm 1, \pm 1)$. Although ϕ might have a non-trivial kernel, the following lemma shows that it does not have a $\mathrm{Gal}_{\mathbb{Q}}$-stable subset.

Lemma 4.9. *Let $x \in X^*(\widetilde{T}')$. If $\phi(\tau(x)) = 0$ for all $\tau \in \mathrm{Gal}_{\mathbb{Q}}$, then $x = 0$.*

Proof. Since the subgroup of $X^*(\widetilde{T}')$ generated by $\{(\pm 1, \cdots, \pm 1)\}$ is of finite index, we may assume without loss of generality that x lies in the subgroup. As τ runs through $\mathrm{Gal}_{\mathbb{Q}}$, $\phi(\tau(x))$ runs through all p-adic valuations of $\mathrm{ev}'(x)$. So these are all 0 and hence, all absolute values of $\mathrm{ev}'(x)$ are 0. So $\mathrm{ev}'(x)$ is a root of unity, which implies that $\mathrm{ev}'(x) = 1$. Since ev' is injective, this is a contradiction. □

First, note that $\mathbb{Z}^5/\ker(\mathrm{ev})$ is torsion-free since the image of ev contains no roots of unity other than 1. Now $\ker(\mathrm{ev})$ is stable under the $\mathrm{Gal}_{\mathbb{Q}_l}$-action and the image of $\mathrm{Gal}_{\mathbb{Q}_l}$ in $\{\pm 1\} \rtimes S_5$ contains a 5-cycle. So $\ker(\mathrm{ev})$ is a $\mathrm{Gal}_{\mathbb{Q}_l}$-stable sublattice of $\mathbb{Z}^5$ with the quotient $\mathbb{Z}^5/\ker(\mathrm{ev})$ torsion-free. Since 5 is a prime , the only possibilities for $\ker(\mathrm{ev})$ are the following:

1. $\ker(\mathrm{ev}) = \mathbb{Z}^5$
2. $\ker(\mathrm{ev}) = \{(x_1, \cdots, x_5) : x_1 + \cdots + x_5 = 0\}$
3. $\ker(\mathrm{ev}) = \{(x, \cdots, x) : x \in \mathbb{Z}\}$
4. $\ker(\mathrm{ev}) = 0$

(see the appendix on cyclic lattices for details. Note that in the case of good ordinary reduction - which occurs with density one- only case 4 can occur.)

In **Case 1**, all eigenvalues of Fr_v are $\sqrt{q}$. So all Newton slopes are $\frac{1}{2}$. Hence, A is isogenous to the 16-th power of the supersingular elliptic curve and $\mathrm{End}^0(A_{\overline{k}}) = \mathrm{Mat}_{16}(\mathbb{Q}_{p,\infty})$, where $\mathbb{Q}_{p,\infty}$ is the quaternion algebra over $\mathbb{Q}$ ramified only at p and ∞. Clearly, $\mathrm{Gal}(\mathbb{Q}(\mathcal{W}_{A_v})/\mathbb{Q}) = \{1\}$.

In **Case 2**, $X^*(\widetilde{T}')$ is a free $\mathbb{Z}$-module of rank 1. Since none of the weights $(\pm 1, \pm 1, \pm 1, \pm 1, \pm 1)$ are in the kernel of ev, none of the simple components correspond to $\sqrt{q}$.

Write $a_1 = (1,1,1,1,1)', a_2 = (-1,1,1,1,1)', \cdots, a_6 = (-1,-1,-1,-1,-1)' \in X^*(\widetilde{T}')$. Then

$$a_1 - a_2 = a_2 - a_3 = a_3 - a_4 = a_4 - a_5 = a_5 - a_6$$

and $\phi(a_1) = -\phi(a_6)$. Since the only conjugates of $\mathrm{ev}(a_1)$ are $\mathrm{ev}(a_1)$ and $\overline{\mathrm{ev}(a_1)}$, lemma 4.9 implies that $\phi(a_1) \neq 0$. Hence, $\phi(a_1) = 5\lambda$, $\phi(a_2) = 3\lambda$, $\phi(a_3) = \lambda$, $\phi(a_4) = -\lambda$, $\phi(a_5) = -3\lambda$, $\phi(a_6) = -5\lambda$ for some λ. So the Newton slopes are $\frac{1}{2} \pm 5\lambda$, $5 \times \frac{1}{2} \pm 3\lambda$, $10 \times \frac{1}{2} \pm \lambda$. Since the slopes lie between 0 and 1 and the Newton polygon has integral break points, it follows that $\lambda = \frac{1}{10}$. Hence, the Newton polygon of A_v is $0, 5 \times \frac{1}{5}, 10 \times \frac{2}{5}, 10 \times \frac{3}{5}, 5 \times \frac{4}{5}, 1$.

Now $\rho_l(\mathrm{Fr}_v)$ has six distinct $\mathrm{ev}(a_1), \cdots, \mathrm{ev}(a_6)$ eigenvalues on V_l. We have $\mathrm{ev}(a_1) = \overline{\mathrm{ev}(a_6)}$ is of degree 2 and hence, $\mathrm{ev}(a_1)\sqrt{q}$ is a Weil q-integer of degree 2. So it corresponds to an ordinary elliptic curve $A^{(1)}$. Write $A_v = A^{(1)} \times A'$. Since A' has Newton polygon $5\times\frac{1}{5}, 10\times\frac{2}{5}, 10\times\frac{3}{5}, 5\times\frac{4}{5}$, every simple component has dimension divisible by 5. Since $\mathrm{Gal}(\mathbb{Q}(\mathcal{W}_{A_v})/\mathbb{Q}) = \{\pm 1\} \times H$ for some $H \subseteq S_5$, it follows that A' has a simple component of dimension 5. So $\mathrm{Gal}_{\mathbb{Q}}$ has precisely three orbits in $\mathcal{W}_{A_v}$ and hence, A' has two simple components (possibly with multiplicities greater than one).

Now suppose, by way of contradiction, that A' has a simple component B of dimension 10. Its Weil number is of degree 2 over $\mathbb{Q}$ and hence, $20 = 2[\mathrm{End}^0(B) : \mathrm{Cent}(\mathrm{End}^0(B))]^{1/2}$. So $[\mathrm{End}^0(B) : \mathrm{Cent}(\mathrm{End}^0(B))]^{1/2} = 10$. But the exact sequence

$$0 \longrightarrow \mathrm{Br}(L|K) \xrightarrow{i} \bigoplus_w \mathrm{Br}(L^w|K_w) \xrightarrow{\alpha \mapsto \sum_v \mathrm{inv}_v(\alpha)} \frac{1}{\mathrm{lcm}(\{n_w\}_w)}\mathbb{Z}/\mathbb{Z} \longrightarrow 0$$

$\big($where L^w is the completion of L at any prime lying over w and $n_w = [L^w : K_w]\big)$ for Galois extensions L/K of global fields implies that $[\mathrm{End}^0(B) : \mathrm{Cent}(\mathrm{End}^0(B))]^{1/2}$ is the least common denominator of the local invariants $\mathrm{inv}_w(\mathrm{End}^0(B))$. But this integer, in turn, divides the least common denominator of the Newton slopes, which is 5. This yields a contradiction.

Thus, A' has two simple components B_1 and B_2 each of dimension 5 and of multiplicities 1 and 2 respectively. Furthermore, both $\mathrm{End}^0(B_1)$ and $\mathrm{End}^0(B_1)$ are 25 dimensional central division algebras over imaginary quadratic fields in which p splits.

In **Case 3**, $X^*(\widetilde{T'})$ is a free $\mathbb{Z}$-module of rank 4. Now $(1,1,1,1,1)' = (-1,-1,-1,-1,-1)' = 0$. The Weil q-integer $\sqrt{q} = \mathrm{ev}(1,1,1,1,1)\sqrt{q}$ corresponds to a supersingular elliptic curve $A^{(1)}$ over k. Furthermore, the other 30 eigenvalues $\mathcal{W}_{A_v} \setminus \mathrm{ev}(\pm(1,1,1,1,1))$ are all distinct. So the endomorphism algebra of the 15 dimensional complement A' of $A^{(1)}$ is a CM algebra of dimension 30.

The image of $\mathrm{Gal}_{\mathbb{Q}}$ in $\{\pm 1\} \rtimes S_5$ contains a cycle of length 5 which induces a cyclic permutation of $\{\mathrm{ev}(-1,1,1,1,1), \cdots, \mathrm{ev}(1,1,1,1,-1)\}$. Since complex conjugation acts by inversion, $\mathrm{Gal}_{\mathbb{Q}}$ acts transitively on $\{\mathrm{ev}(\pm(-1,1,1,1,1)), \cdots, \mathrm{ev}(\pm(1,1,1,1,-1))\}$. Similarly, $\mathrm{Gal}_{\mathbb{Q}}$ acts transitively on

$$\{\mathrm{ev}(\pm(-1,-1,1,1,1)), \cdots, \mathrm{ev}(\pm(-1,1,1,1,-1))\}$$

and

$$\{\mathrm{ev}(\pm(-1,1,-1,1,1)), \cdots, \mathrm{ev}(\pm(1,-1,1,1,-1))\}.$$

Set $\pi_1 = \mathrm{ev}(-1,1,1,1,1)\sqrt{q}$, $\pi_2 = \mathrm{ev}(-1,-1,1,1,1)\sqrt{q}$, $\pi_3 = \mathrm{ev}(-1,1,-1,1,1)\sqrt{q}$. Each π_i $(i = 1,2,3)$ is a Weil q-integer of degree a multiple of 10. Let A_i be the corresponding simple abelian variety over $\mathbb{F}_q$ (upto isogeny), L_i the field $\mathbb{Q}(\pi_i)$ and $\alpha_i = \frac{\pi_i}{\sqrt{q}}$. Since $\mathrm{End}^0(A_i) = L_i$, it follows that the local invariants $[L_{i,v} : \mathbb{Q}_p]v(\tau(\pi_i)) \in \mathbb{Z}$ for all $\tau \in \mathrm{Gal}_{\mathbb{Q}}$.

Furthermore, we have $\Phi_{\pi_1} = \Phi_{\pi_2} = \Phi_{\pi_3}$ and $\Phi_{\pi_0} = \langle\sqrt{q}\rangle$. Now, $\mathrm{Gal}(\mathbb{Q}(\mathcal{W}_{A_v})^+/\mathbb{Q})$ is a transitive subgroup of S_5. So it must be one of the following groups:

$$\{\langle(1\ 2\ 3\ 4\ 5)\rangle, \langle(1\ 2\ 3\ 4\ 5), (1\ 5)(2\ 4)\rangle, \langle(1\ 2\ 3\ 4\ 5), (1\ 2\ 5\ 4)\rangle, A_5, S_5\}.$$

So every simple component of A' has dimension a multiple of 5. Suppose by way of contradiction that every simple component is a five-fold. Then, in particular, π_2 is not conjugate to π_3. Hence, $\mathrm{Gal}(\mathbb{Q}(\mathcal{W}_{A_v})^+/\mathbb{Q})$ is cyclic of order 5 and $\mathrm{Gal}(\mathbb{Q}(\mathcal{W}_{A_v})/\mathbb{Q})$ is cyclic of order 10. Then $\mathrm{End}^0(A')$

is commutative and it follows from lemma 4.6 that every simple component of A' is ordinary. Hence, the Newton polygon is $2 \times \frac{1}{2}, 15 \times 0, 15 \times 1$ which is not possible as noted in subsection 4.3. So $\mathrm{Gal}(\mathbb{Q}(\mathcal{W}_{A_v})^+/\mathbb{Q})$ is one of the other four possibilities and $\pi_2 = \mathrm{ev}(-1,-1,1,1,1)\sqrt{q}$ is conjugate to $\pi_3 = \mathrm{ev}(-1,1,-1,1,1)\sqrt{q}$. Hence, the simple component corresponding to π_2 is of dimension 10.

The endomorphism algebra $\mathrm{End}^0(A_v) \cong \mathbb{Q}_{p,\infty} \times F_{10} \times F_{20}$ where the subscript is the degree of the CM field and $F_{10} \hookrightarrow F_{20}$.

In **Case 4**, there are 32 distinct eigenvalues and the endomorphism algebra is a CM algebra of degree 32. An argument identical to the one in case 3 shows that $\mathrm{Gal}_{\mathbb{Q}}$ acts transitively on the sets $\{\mathrm{ev}(\pm(1,1,1,1,1))\}$, $\{\mathrm{ev}(\pm(-1,1,1,1,1)), \cdots, \mathrm{ev}(\pm(1,1,1,1,-1))\}$, $\{\mathrm{ev}(\pm(-1,-1,1,1,1)), \cdots, \mathrm{ev}(\pm(-1,1,1,1,-1))\}$ and $\{\mathrm{ev}(\pm(-1,1,-1,1,1)), \cdots, \mathrm{ev}(\pm(1,-1,1,1,-1))\}$. So the simple component corresponding to the Weil number π_0 has dimension $\equiv 1 \pmod 5$ and the other simple components have dimensions divisible by 5. Furthermore, we have $\mathrm{Gal}(\mathbb{Q}(\mathcal{W}_{A_v})/\mathbb{Q}) \subseteq \{\pm 1\}^j \rtimes H$ and since H fixes $\pi_0 = \mathrm{ev}(1,1,1,1,1)\sqrt{q}$, we have $\mathbb{Q}(\pi_0) \subseteq \mathbb{Q}(\mathcal{W}_{A_v})^H$. So $[\mathbb{Q}(\pi_0) : \mathbb{Q}] = 2^i$ for some $1 \leq j \leq 5$ and $2^{i-1} \equiv 1 \pmod 5$, which implies that $i = 1$ or 5.

Also, we have

$$\Phi_{\pi_0} \subseteq \Phi_{\pi_1} \subseteq \Phi_{\pi_2} = \Phi_{\pi_3} = \Phi_{A_v}.$$

Furthermore, we have the inclusion $\mathbb{Q}(\pi_0)^{\mathrm{Gal}} \subseteq \mathbb{Q}(\pi_1)^{\mathrm{Gal}} = \mathbb{Q}(\pi_2)^{\mathrm{Gal}} = \mathbb{Q}(\pi_3)^{\mathrm{Gal}}$ of Galois closures and $[\mathbb{Q}(\pi_0) : \mathbb{Q}] \geq 2$, $[\mathbb{Q}(\pi_1) : \mathbb{Q}] \geq 10$, $[\mathbb{Q}(\pi_2) : \mathbb{Q}] \geq 10$, $[\mathbb{Q}(\pi_3) : \mathbb{Q}] \geq 10$. Thus, the special fiber A_v has one of the following simple decompositions in **Case 4**:

(i) Simple of dimension 16

In this case, we have $[\mathbb{Q}(\pi_0) : \mathbb{Q}] = 32$. Write $\mathrm{Gal}(\mathbb{Q}(\mathcal{W}_{A_v})/\mathbb{Q}) = \{\pm 1\}^5 \rtimes H$. Since H fixes π_0, it follows that A_v has CM by the field $\mathrm{End}^0(A_v) \cong \mathbb{Q}(\pi_0)$, the subfield of $\mathrm{Gal}(\mathbb{Q}(\mathcal{W}_{A_v})/\mathbb{Q})$ fixed by H.

In all other cases, π_0 corresponds to an ordinary elliptic curve. Suppose, by way of contradiction, that A_v is not simple and $\mathrm{Gal}(\mathbb{Q}(\mathcal{W}_{A_v})/\mathbb{Q}(\mathcal{W}_{A_v})^+)$ is of rank 5. Choose an element $\tau_j \in \mathrm{Gal}(\mathbb{Q}(\mathcal{W}_{A_v})/\mathbb{Q}(\mathcal{W}_{A_v})^+)$ that fixes $\sigma^j(\pi_2)$ and swaps $\sigma^i(\pi_2)$ and $\sigma^j(\overline{\pi}_2)$ for $i \neq j$, $(0 \leq i, j \leq 4)$.

We have

$$\pi_0 = \pi_2 \sigma(\overline{\pi}_2)^{-1} \sigma^2(\pi_2) \sigma^3(\overline{\pi}_2)^{-1} \sigma^4(\pi_2).$$

Then $\pi_0 \pi_0^{\tau_0} = \pi_2^2$. So $\pi_0 \pi_0^{\tau_0} = q$ or π_0^2 depending on whether τ_0 fixes π_0. So either $\pi_2^2 = q$ or $\pi_2^2 = \pi_0^2$, neither of which is possible in **Case 4**.

Let A' be the 15-dimensional complement of A_{π_0} in A_v and let B be an odd dimensional simple component of A'. Let π_i be the corresponding Weil number (where $i = 1$, 2 or 3). Then $\mathbb{Q}(\pi_i + \overline{\pi}_i)$ is of odd degree over $\mathbb{Q}$ and hence, lies in the fixed field of $\{\pm 1\}$. Since $\{\pm 1\}$ is a normal subgroup of $\mathrm{Gal}(\mathbb{Q}(\mathcal{W}_{A_v})/\mathbb{Q})$, the Galois closure $\mathbb{Q}(\mathcal{W}_{A_v})^+$ of $\mathbb{Q}(\pi_i + \overline{\pi}_i)$ also lies in the fixed field of $\{\pm 1\}$. Hence, $\mathrm{Gal}(\mathbb{Q}(\mathcal{W}_{A_v})/\mathbb{Q}) = \{\pm 1\} \times \mathrm{Gal}(\mathbb{Q}(\mathcal{W}_{A_v})^+/\mathbb{Q})$.

(ii) $A_v = A^{(1)} \times A^{(5)} \times A^{(10)}$

In this case, π_2, π_3 lie in the same $\mathrm{Gal}_{\mathbb{Q}}$-conjugacy class. So $\mathrm{Gal}(\mathbb{Q}(\mathcal{W}_{A_v})/\mathbb{Q}) = \{\pm 1\} \times H$ where H is a transitive subgroup of S_5 of order divisible by 10.

(iii) $A_v = A^{(1)} \times A_1^{(5)} \times A_2^{(5)} \times A_3^{(5)}$

In this case, $\mathbb{Q}(\pi_0)$ is imaginary quadratic with p split in $\mathbb{Q}(\pi_0)/\mathbb{Q}$ and the elements π_i $(i = 1, 2, 3)$

are pairwise non-conjugate and of degree 10. Since π_2, π_3 are non-conjugate, it follows that $\mathrm{Gal}(\mathbb{Q}(\mathcal{W}_{A_v})^+/\mathbb{Q})$ is cyclic of order 5 and $\mathrm{Gal}(\mathbb{Q}(\mathcal{W}_{A_v})^+/\mathbb{Q})$ is cyclic of order 10. So p splits completely in $\mathbb{Q}(\mathcal{W}_{A_v})/\mathbb{Q}$ and hence, by lemma 4.7, every simple component is ordinary.

(iv)' $A_v = A^{(1)} \times A^{(15)}$
In this case, π_1, π_2, π_3 lie in the same conjugacy class and π_0 is of degree 2. Furthermore, 30 divides $|\mathrm{Gal}(\mathbb{Q}(\mathcal{W}_{A_v})/\mathbb{Q})|$ and hence, $\mathrm{Gal}(\mathbb{Q}(\mathcal{W}_{A_v})/\mathbb{Q}) = \{\pm 1\}^j \rtimes H$ with $j = 1$ or 5 and $H = A_5$ or S_5. Since $\{\sigma^i(\pi_1), \sigma^i(\overline{\pi}_1) : 1 \leq i \leq 5\}$ is stable under the action of H and the complex conjugation, it follows that $j = 5$, a contradiction. □

We briefly describe how the techniques in the preceding theorem may be used for higher dimensions. Consider an abelian variety A weakly of Mumford type of rank $2n+2$ with $2n+1$ a prime. As explained in appendix 6.2, ker(ev) has the same four possibilities as above.

Clearly, if ker(ev) is of rank $2n+1$, then all the eigenvalues are $\frac{1}{2}$ and the reduction is isogenous to the 2^{2n}-th power of the supersingular elliptic curve.

If ker(ev) is the index 1 sublattice $\{(x_1, x_2, \cdots, x_N) : \sum_{i=1}^{N} x_i = 0\}$, then A_v has precisely $2n+2$ distinct eigenvalues. Furthermore, with techniques identical to those in the preceding theorem, we may show that the Newton polygon of A_v is

$$2^{2n+1-j}\binom{2n+1}{j} \times \frac{j}{2n+1}, \quad 0 \leq j \leq 2n+1.$$

So every simple component of A_v other than the ordinary elliptic curve has dimension a multiple of $2n+1$. Furthermore, for any such simple component B, the integer $[\mathrm{End}^0(B) : \mathrm{Cent}(\mathrm{End}^0(B))]^{1/2}$ is the least common multiple of the Hasse invariants $\mathrm{inv}_w([\mathrm{End}^0(B)])$ and hence, must divide $2n+1$. Since $2n+1$ is a prime by assumption, it follows that $[\mathrm{End}^0(B) : \mathrm{Cent}(\mathrm{End}^0(B))] = (2n+1)^2$. So each $\mathrm{End}^0(B)$ is a $(2n+1)^2$-dimensional central division algebra over an imaginary quadratic field in which p splits. Thus, by Honda-Tate theory, every simple component other than the ordinary elliptic curve has dimension $2n+1$ and $\mathrm{Gal}(\mathbb{Q}(\mathcal{W}_{A_v})/\mathbb{Q})$ is abelian of exponent 2. The simple component of A_v corresponding to the Weil number $\mathrm{ev}(\underbrace{-1, \cdots, -1}_{j}, 1, \cdots, 1)\sqrt{q}$ occurs with multiplicity $\frac{1}{2n+1}\binom{2n+1}{j}$ and has precisely two Newton slopes $\frac{j}{2n+1}, \frac{2n+1-j}{2n+1}$, each with multiplicity $2n+1$.

If $\ker(\mathrm{ev}) = \{(x, x, \cdots, x) : x \in \mathbb{Z}\}$, then $\mathrm{Gal}(\mathbb{Q}(\mathcal{W}_{A_v})/\mathbb{Q}) = \{\pm 1\} \times H$ where H is some subgroup of S_{2n+1} containing a $2n+1$-cycle. Clearly, H determines the $\mathrm{Gal}_{\mathbb{Q}}$-orbits of $\mathcal{W}_{A_v} \setminus \{\sqrt{q}\}$ and hence, determines the simple decomposition of the complement of the supersingular elliptic curve in A_v.

We now state a result in the case when ker(ev) is trivial. As before, we denote the Weil number $\mathrm{ev}(1, \cdots, 1)\sqrt{q}$ by π_0.

Proposition 4.10. *Let $2n+1$ be a prime. Let A be an abelian variety over a number field, weakly of Mumford type of rank $2n+2$ and let v be a finite place of F. Then $[\mathbb{Q}(\pi_0) : \mathbb{Q}] = 2^i$ for some $1 \leq i \leq 2n+1$ such that $2^i \equiv 1 \pmod{2n+1}$ and A_v is simple if and only if $[\mathbb{Q}(\pi_0) : \mathbb{Q}] = 2^{2n+1}$.*

Proof. By lemma 3.5, we may enlarge the field if necessary to ensure good reduction at v. Furthermore, the kernel of the map $\mathrm{ev} : X^*(T) \longrightarrow \Phi_{A_v}$ has the four possibilities listed in appendix 6.2. Unless $\ker(\mathrm{ev}) = 0$, we see that π_0 corresponds to an elliptic curve. So we may

assume without loss of generality that $\ker(\mathrm{ev}) = 0$. So $P_{A_v}(X)$ is separable and $\mathrm{End}^0(A_v)$ is a CM algebra of degree 2^{2n+1}.

Now, note that $\mathrm{Gal}(\mathbb{Q}(\mathcal{W}_{A_v})/\mathbb{Q}) = \{\pm 1\}^j \rtimes H$ where some $1 \leq j \leq 2n+1$ and H is a transitive subgroup of S_{2n+1}. Since π_0 is fixed by H, we have $\mathbb{Q}(\pi_0) \subseteq \mathbb{Q}(\mathcal{W}_{A_v})^H$ and since $[\mathbb{Q}(\mathcal{W}_{A_v})^H : \mathbb{Q}] = 2^j$, it follows that $[\mathbb{Q}(\mathcal{W}_{A_v})^H : \mathbb{Q}] = 2^i$ for some $i \leq j$. Furthermore, since $2n+1$ is assumed to be a prime, H contains a $2n+1$-cycle. Hence, every simple component of A_v corresponding to a Weil number non-conjugate to π_0 has dimension a multiple of $2n+1$ and the component corresponding to π_0 has dimension $\equiv 1 \pmod{2n+1}$.

Since the eigenvalues are distinct, it is clear that A_v is simple if and only if $[\mathbb{Q}(\pi_0) : \mathbb{Q}] = 2 \dim A_v = 2^{2n+1}$. □

In particular, we note that at places where the abelian variety has simple reduction, the eigenvalues of the Frobenius are all distinct and the endomorphism ring is an order in a CM field. So the image of Fr_v in $\mathrm{End}(T_l(A))$ generates a maximal torus of the image of the Galois representation ρ_l.

4.5 Endomorphism Algebras

We now list the possibilities for $\mathrm{End}^0(A_{\overline{k}})$ that we derived in the course of the last proof. Given the division algebra $\mathrm{End}^0(Y)$ with center C for a simple component Y, we can easily retrieve the dimension through the equality $\dim Y = \frac{1}{2}[\mathrm{End}^0(Y) : C][C : \mathbb{Q}]$. Hence, this is a more concrete characterization that the dimensions of the simple components, even though the endomorphism algebras do not determine the precise isogeny classes for abelian varieties of genus greater than one.

We introduce some notations for brevity. The center of the endomorphism algebra of a simple component of $A_{\overline{k}}$ is some field $\mathbb{Q}(\sqrt{q}\alpha)$ with $\sqrt{q}\alpha$ a Weil q-integer. So it is either $\mathbb{Q}$ (which only happens in the supersingular case) or a CM field. We use the symbol $\mathbb{Q}_{p,\infty}$ for the unique quaternion algebra over $\mathbb{Q}$ ramified ony at p and ∞.

We will use F_{2i} to mean a CM field of degree $2i$ and $D^{2i,j}$ to mean the central division algebra of dimension j^2 over a CM field E_{2i} of degree $2i$ such that:
- the rational prime p splits completely in E_{2i};
- $D^{2i,j}$ has Hasse invariants

$$\mathrm{inv}_w(D^{2i,j}) = \begin{cases} 0 & \text{if } w \nmid p \\ \frac{w(\pi)}{w(q)}[E_{2i,w} : \mathbb{Q}_p] \pmod 1 & \text{if } w|p. \end{cases}$$

Proposition 4.11. *For an abelian variety A weakly of Mumford type of rank 6 over a number field F, $\mathrm{End}^0(A_{\overline{k}})$ has the following possibilities:*

(1) $\mathrm{Mat}_{16}(\mathbb{Q}_{p,\infty})$ (*Supersingular case*)
(2) $F_2 \times D_1^{2,5} \times \mathrm{Mat}_2(D_2^{2,5})$
(3) $\mathbb{Q}_{p,\infty} \times F_{10} \times F_{20}$
(4) $F_2 \times F_{10} \times F_{20}$
(5) $F_2 \times F_{10} \times F_{10} \times F_{10}$ *with* $\mathrm{Gal}(F_{10}/\mathbb{Q}) = \mathbb{Z}/10\mathbb{Z}$, $F_2 \subseteq F_{10}$ *and* $p \in \mathcal{S}(F_{10}|\mathbb{Q})$
(6) F_{32}

The last possibility occurs with Dirichlet density one.

Proof. This is shown in the proof of the preceding theorem. The density of places with simple ordinary reduction will be shown in section 5. □

Remark We note that in the cases where the endomorphism algebra is a CM algebra, the equality of endomorphism algebras holds over the smallest extension of k over which the reduction is well-defined.

4.6 An existence result in the ordinary case

In the preceding subsection, we saw that for an abelian variety weakly of Mumford type of rank 6, the reduction at a place of good ordinary reduction is either simple or of the form $A^{(1)} \times A'$, where A is an elliptic curve. In this section, we show the existence of the latter possibility. Except for Proposition 4.13, the techniques are virtually identical to those from [No01] and hence, we merely provide a sketch.

The tori T'_L and T_L. We introduce some notation here. Let L be a CM field and let L^+ be its totally real subfield of index 2. The norm map $N_{L/L^+} : L^* \longrightarrow L^{+*}$ defines a map $\mathrm{Res}_{L/\mathbb{Q}}(\mathbb{G}_m) \longrightarrow \mathrm{Res}_{L^+/\mathbb{Q}}(\mathbb{G}_m)$. We define T'_L to be the kernel of the map and T_L to be the pre-image of $\mathbb{G}_{m,\mathbb{Q}}$.

Now let K be a totally real field of degree $2n+1$, with $2n+1$ a prime. For simplicity, we assume K is Galois over $\mathbb{Q}$. Let D be a quaternion over K split at exactly one real place and with trivial corestriction in $\mathrm{Br}(\mathbb{Q})$. By Skolem-Noether, we have $(D^*)^{\mathrm{der}} = \{d \in D^* : d\overline{d} = 1\}$. The datum $((D^*)^{\mathrm{der}}, h)$ gives a Shimura curve that is of Hodge type but not of PEL type. We refer the reader to ([SZ12], Section 2) for more details.

Let X be an abelian variety strongly of Mumford type that is a generic point on the Shimura curve given by $((D^*)^{\mathrm{der}}, h)$. It has a model over the field K ([No01], [SZ12]). Let v be a place of K of good reduction. Since D is ramified at $2n$ of the archimedean places, any maximal subfield of D is necessarily imaginary quadratic field over K. The following result of Tate guarantees the existence of an intermediate CM field between a totally real field and a quaternion algebra over it.

Proposition 4.12. (Tate) *Let K be a totally real field and let D be a d^2-dimensional central division algebra over K. If d is even, there exists a CM field L with an embedding in D such that $[L : K] = d$.*

A proof may be found in ([CCO09], Chapter 1). We will need the following variant for our purposes, a proof for which we give in appendix 6.3.

Proposition 4.13. *Let K be a (totally) real field Galois over $\mathbb{Q}$ and a D a $4d^2$-dimensional central division algebra over K. If $2 \nmid d$ and $\gcd(2d, [K : \mathbb{Q}]) = 1$, there exists a totally imaginary number field E cyclic of degree $2d$ over $\mathbb{Q}$ such that $[EK : K] = 2d$ and $EK \hookrightarrow D$.*

In particular, there exists an imaginary quadratic field E such that $L = EK$. The inclusion $L^* \subseteq D^*$ induces morphisms $\tilde{\rho}' : T'_L \longrightarrow \tilde{G}'$ and $\rho' = N \circ \tilde{\rho}' : T'_L \longrightarrow G'$. Let $\phi_1, \cdots, \phi_{2n+1}, \overline{\phi}_1, \cdots, \overline{\phi}_{2n+1}$ be the embeddings $L \hookrightarrow \mathbb{C}$. Let ϕ_{2n+1} be the embedding corresponding to the archimedean place where D is split. We write $[\phi_j]$, $[\overline{\phi_j}]$ for the induced characters of L^*, of T^*_L and of T'_L respectively. So $[\phi_1], \cdots, [\phi_{2n+1}]$ is a basis for $X^*(T'_L)$. We fix an isomorphism $L \otimes_{\mathbb{Q}} \mathbb{R} \cong \mathbb{C}^{2n+1}$ such that the composite of $L \hookrightarrow L \otimes \mathbb{R}$ with the projection on the i-th factor $\mathbb{C}$ induces the embedding $\phi_i : L \hookrightarrow \mathbb{R}$. We deduce an isomorphism $L^*_{\mathbb{R}} \cong S^{2n+1}$. Define h to be the composite

$$S \longrightarrow L^*_{\mathbb{R}} \hookrightarrow D^*_{\mathbb{R}} \longrightarrow G_{\mathbb{R}}$$

where the first map is the inclusion of the coordinate corresponding to ϕ_{2n+1} and the last map is the norm.

For $h' = h|_{S^1} \longrightarrow G'_{\mathbb{R}}$, there is a map $\tilde{h'} : S^1 \longrightarrow (T'_L)_{\mathbb{R}} \subseteq G'_{\mathbb{R}}$ such that $h' = \rho' \circ \tilde{h'}$. Now h is conjugate to the map h_0 and factors through the torus $T \subseteq G$, the image of $L^* \subseteq D^*$. So h defines a set of special points of the Mumford curve. Let X be an abelian variety in the corresponding isogeny class. Then X is of dimension 2^{2n} and has CM. At any place v of good ordinary reduction, we have $\mathrm{End}^0(X) = \mathrm{End}^0(X_v)$. So to study the splitting of X_v, it suffices to study that of X.

Proposition 4.14. *With the setup as above, X is isogenous to a product $X^{(1)} \times X'$ where $X^{(1)}$ is an ordinary elliptic curve with CM by the field E.*

Proof. Write $2^{2n} - 1 = m(2n+1)$. We define the map $\tau : T'_L \longrightarrow T'_L$ by $x \mapsto N_{L/E}(x)x^{-2}$. This yields the map

$$\pi' = (\tau^{(m)}, N_{L/E}) : T'_L \longrightarrow T_L'^{(m)} \times T'_E.$$

The natural action of $T_L'^{(m)} \times T'_E$ on $L^{(m)} \oplus E \cong \mathbb{Q}^{2^{2n+1}}$ makes π' into a 2^{2n+1}-dimensional representation of T'_L. The weights of the representation induced by $N_{L/E}$ are $\pm\left(([\phi_1] + \cdots + [\phi_{2n+1}])\right)$ and those of the representation induced by τ are $\sum_{i\in\mathcal{I}}[\phi_i] - \sum_{j\in\mathcal{J}}[\phi_j]$, where $\mathcal{I}, \mathcal{J}$ run through the pairs of non-empty sets $\mathcal{I}, \mathcal{J}$ such that $\mathcal{I} \cup \mathcal{J} = \{1, \cdots, 2n+1\}$, $\mathcal{I} \cap \mathcal{J} = \emptyset$. So the weights of π' are $\sum_{i=1}^{2n+1} \pm[\phi_i]$. These coincide with the weights of the representation $\rho' = N \circ \tilde{\rho'} : T'_L \to G'$ and hence, $\pi' = \rho'$. Thus, X_v is isogenous to a product $X^{(1)} \times X'$ where $X^{(1)}$ is an ordinary elliptic curve with CM by E and X' is an abelian variety (not necessarily simple) with an action of L. □

Remark The $2^{2n} - 1$-dimensional complement X' is not simple. The Galois group $\mathrm{Gal}(\mathcal{W}_{X_v}/\mathbb{Q})$ is $\{\pm 1\} \times H$ for some transitive $H \subseteq S_{2n+1}$ and hence, the simple decomposition of X' is determined by H.

The following Proposition now implies the existence of generic points on the Mumford curve such that the corresponding abelian variety has a reduction at some prime that has an ordinary elliptic curve as a simple component.

Proposition 4.15. *With the setup as above, let x_F be a special point on the Mumford curve $\mathcal{M}/F$ and X the corresponding abelian variety with complex multiplication. Let $\mathfrak{p}$ be a prime of F with residue field k. There exists a finite extension F' of F, a prime $\mathfrak{p}'$ lying over $\mathfrak{p}$ and a generic point $y_{F'} \in \mathcal{M}(F')$ such that the abelian variety Y corresponding to $y_{F'}$ has reduction $Y_{k'}$ isogenous to $X_{k'}$.*

The proof is identical to that of ([No01], Proposition 5.1) and hence, is omitted.

We conclude this section by briefly describing the ordinary elliptic curves over finite fields that occur as simple components of the reductions X_v. As seen above, the elliptic curve $X^{(1)}$ has CM by the imaginary quadratic field E. Since $\mathrm{End}^0(X^{(1)})$ determines $X^{(1)}$ upto isogeny, it suffices to determine the possibilities for the endomorphism algebra $\mathrm{End}^0(X^{(1)})$.

Let $\mathfrak{l}_1, \cdots, \mathfrak{l}_t$ be the finite prime of K and let l_i be the rational prime lying under $\mathfrak{l}_i$. Now $EK \hookrightarrow D$ if and only if EK splits D. Since EK is a CM field, this is equivalent to EK having

local degree 2 at all the finite primes of K at every $\mathfrak{l}_i$, $1 \leq i \leq t$. But since we assumed that $K/\mathbb{Q}$ is Galois of odd degree, this is equivalent to $E/\mathbb{Q}$ being either ramified or inert at each of the rational primes l_i.

5 Density of places of simple reduction

We now look into the density of places where the different types of reductions occur. Since the abelian varieties weakly of Mumford type have trivial endomorphism rings, the following conjecture suggests that they have absolutely simple reduction at a set of places of density one.

Conjecture. Let X be an abelian variety over a number field F such that $F_X^{\mathrm{conn}} = F$, and $\mathrm{End}(X_{\overline{\mathbb{Q}}})$ is commutative. Then X has absolutely simple reduction away from a set of places of density zero.

While the conjecture remains open in this generality, several special cases have been studied. If the abelian variety is, in fact, *strongly* of Mumford type, the conjecture holds by the following theorem from [Zyw14].

Theorem 5.1. [Zyw14] *Let X be an abelian variety over a number field F such that $F^{\mathrm{conn}} = F$. If $\mathrm{End}^0(X_{\overline{\mathbb{Q}}})$ is a commutative field and the Mumford-Tate conjecture holds for X, then X has absolutely simple reduction away from a density zero set of places of F.*

From this, one can easily deduce the following Proposition.

Proposition 5.2. *Let A be an abelian variety strongly of Mumford type over a number field F such that $F_A^{\mathrm{conn}} = F$. Then A has absolutely simple reduction at a set of places of density one.*

Proof. Since the root system of $\mathrm{Lie}(\mathrm{MT}(A))$ is of type A_1, the Mumford-Tate conjecture holds by Theorem 5.13 in [Pin98]. Since $\mathrm{End}(A_{\overline{\mathbb{Q}}}) = \mathbb{Z}$, this result now follows immediately from Zywina's theorem. □

We now study the reductions of abelian varieties weakly of Mumford type. While the Mumford-Tate conjecture remains open for this class for rank greater than 4 (alternatively dimension greater than 4), we know from [Pin98, 5.13] that there exists a quasi-split reductive group that looks like the Mumford-Tate group.

5.1 Reductions of abelian varieties weakly of Mumford type

Throughout the rest of this section, unless otherwise specified, let A be an abelian variety of rank $2n+2$ over a number field F, weakly of Mumford type. Some of the results generalize to absolutely simple abelian varieties with trivial endomorphism rings. Enlarging F if necessary, we assume all the G_l are connected. By Theorem 5.11 of [Pin98], there exists a quasi-split connected reductive group $\mathbf{G}_A$ over $\mathbb{Q}$ with a faithful $\mathbb{Q}$-linear representation $\mathbf{V}_A$ such that

-$(\mathbf{G}_A, \mathbf{V}_A) \otimes \mathbb{Q}_l \cong (G_l, V_l)$ for all primes l in a set $\mathbb{S}_A$ of density one.

-$\mathbf{G}_A^{\mathrm{der}}$ is $\mathbb{Q}$-simple.

This section is heavily reliant on adapting the techniques from [Zyw14] and [No09]. To avoid confusion with the notation, we emphasize that while in [Zyw14] the group $\mathbf{G}_A$ is the Mumford-Tate group of A, we use the symbol for the reductive group over $\mathbb{Q}$ described above. The two groups are the same if we assume the Mumford-Tate conjecture for A (which remains open for our class of abelian varieties, as far as we know).

We define the set $\mathcal{S}_A$ of the places v of F such that:

1. A has good ordinary reduction at v.
2. v has local degree 1 over $\mathbb{Q}$.
3. Φ_{A_v} is a free abelian group with rank $\Phi_{A_v} = \text{rank}(A)$.

The first condition holds a set of places of density one by ([Pin98], Theorem 7.13). The second holds for a set of places of density one by the Chebotarev density theorem. That the third condition holds for a density one set is an implication of the following result from [LP97].

For every place v, fix a semisimple element t_v in the conjugacy class of $\rho_l(\text{Fr}_v)$ and let $\mathbf{T}_{v,l}$ be its Zariski closure in G_l.

Theorem 5.3. ([LP97], Theorem 1.2) *For any l, there is a Zariski closed subvariety $Y \subseteq G_l$ such that $\mathbf{T}_{v,l}$ is a maximal torus of G_l whenever $\rho_l(\text{Fr}_v) \in G_l \setminus Y$.*

Now $\mathbf{T}_{v,l}$ is a maximal torus if and only if Φ_{A_v} is a free abelian group of the same rank as the rank of G_l. Thus, the third condition holds for a set of primes of density 1 and hence, $\mathcal{S}_A$ has density 1. So it suffices to show that away from a density zero subset of $\mathcal{S}_A$, A has absolutely simple reduction.

We note that for any $v \in \mathcal{S}_A$ and prime $l \in \mathbb{S}_A$ other than $\text{char}(v)$, the element $\rho_l(\text{Fr}_v)$ is semisimple and hence, lies in some maximal torus of $G_l(\mathbb{Q}_l) = \mathbf{G}_A(\mathbb{Q}_l)$. So for any maximal torus $\mathbf{T}$ of $\mathbf{G}_A$, $\mathbf{T}(\overline{\mathbb{Q}}_l)$ contains an element t conjugate to $\rho_l(\text{Fr}_v)$ in $\mathbf{G}_A(\overline{\mathbb{Q}}_l)$.

The following Proposition shows that good ordinary reduction implies Φ_{A_v} being of full rank if $2n+1$ is a prime and A is an abelian variety weakly of Mumford type of rank $2n+2$.

Proposition 5.4. *Suppose the integer $2n+1$ is a prime. Let v be a place of good ordinary reduction (or more generally, a place such that $\text{End}(A_v)$ is commutative). Then the map $\text{ev} : X^*(\mathbf{T}) \longrightarrow \Phi_{A_v}$ is an isomorphism.*

Proof. Since the map is surjective, it suffices to show injectivity. But since $2n+1$ is a prime, appendix 6.2 shows that $\ker(\text{ev})$ is one of the following:

(a) 0
(b) $\mathbb{Z}^{2n+1}$
(c)$\{(x, \cdots, x) : x \in \mathbb{Z}\}$
(d)$\{(x_1, x_2, \cdots, x_{2n+1} : \sum_{i=1}^{N} x_i = 0\}$

In case (b), A_v is supersingular. In case (c), $\text{ev}(1, \cdots, 1) \in \ker(\text{ev})$ and in case (d), $\rho_l(\text{Fr}_v)$ has precisely $2n+2$ distinct eigenvalues. So in all cases but (a), we see that $P_{A_v}(X)$ is inseparable and $\text{End}(A_v)$ is non-commutative. Hence, ev is injective, which completes the proof. □

Let $\overline{\rho}_l : \text{Gal}_F \longrightarrow \text{Aut}_{\mathbb{Z}/l\mathbb{Z}}(A[l])$ be the representation $\sigma \mapsto (P \mapsto P^\sigma)\ \forall\ P \in A[l]$. Then $\overline{\rho}_l(\text{Gal}_F)$ injects into $\mathcal{G}_l(\mathbb{F}_l)$. The following Proposition shows that it is not much smaller than the full $\mathcal{G}_l(\mathbb{F}_l)$.

Proposition 5.5. [Ser00] *Suppose $F_X^{\text{conn}} = F$ and X is absolutely simple.*
1. *For l sufficiently large, $\mathcal{G}_l$ is a reductive group over $\mathbb{Z}_l$.*
2. *There is a constant C such that $[\mathcal{G}_l(\mathbb{F}_l) : \overline{\rho}_l(\text{Gal}_F)] \leq C$.*
3. *For l sufficiently large, the group $\overline{\rho}_l(\text{Gal}_F)$ contains the commutator subgroup $\mathcal{G}_l(\mathbb{F}_l)$.*

A sketch of the proof may be found in [Zyw14]. More details may be found in [Ser00] and [Win02].

Combining all l-adic representations attached to A, we define the representation

$$\rho_A : \mathrm{Gal}_F \longrightarrow \prod_l \mathrm{Aut}_{\mathbb{Q}_l}(V_l(A)), \quad \sigma \mapsto (\rho_l(\sigma))_l.$$

After enlarging the field F if necessary, the representations ρ_l are independent, by the following result of Serre.

Proposition 5.6. ([Ser00], 138) *There exists a finite extension F'/F such that $\rho_A(\mathrm{Gal}_{F'}) = \prod_l \rho_l(\mathrm{Gal}_{F'})$*

This field extension will be important in measuring the density of the places of F whose Frobenius lands in a fixed conjugacy class of $\overline{\rho}_l(\mathrm{Gal}_F)$.

Proposition 5.7. ([Zyw14], Proposition 2.12) *Let F'/F be an extension as in Serre's lemma. Let Λ be a finite set of rational primes. For each $l \in \Lambda$, fix a subset U_l of $\overline{\rho}_l(\mathrm{Gal}_F)$ stable under conjugation. The set of places v of F such that $\overline{\rho}_l(\mathrm{Fr}_v) \subseteq U_l \; \forall \; l \in \Lambda$ has density*

$$\sum_C \frac{|C|}{\mathrm{Gal}(F'/F)} \prod_{l \in \Lambda} \frac{|\overline{\rho}_l(\Gamma_C) \cap U_l|}{|\overline{\rho}_l(\Gamma_C)|}$$

where C varies over the conjugacy classes of $\mathrm{Gal}(F'/F)$ and $\Gamma_C := \{\sigma \in \mathrm{Gal}_F : \sigma|_{F'} \in C\}$.

5.2 The groups $W(G,T)$ and $\Pi(G,T)$

Let G be a reductive group of a field K. Its maximal tori are all conjugate to each other. Choose a maximal torus T of G. We have a homomorphism

$$\phi_T : \mathrm{Gal}_K \longrightarrow \mathrm{Aut}(X^*(T)), \quad \sigma \mapsto \; (\chi \mapsto \sigma(\chi))$$

for $\sigma \in \mathrm{Gal}_K$, $\chi \in X^*(T)$. Note that T is split if and only if $\phi_T(\mathrm{Gal}_K) = \{1\}$.

We use the symbol $W(G,T)$ to mean the Weyl group $N_G(T)(\overline{K})/T(\overline{K})$ where $N_G(T)$ is the normalizer of T in G. Now, Gal_K acts on $W(G,T)$ and $\sigma(w) = \phi_T(\sigma) \circ w \circ \phi_T(\sigma)^{-1}$ for $\sigma \in \mathrm{Gal}_K$, $w \in W(G,T)$. We define $\Pi(G,T)$ to be the subgroup of $\mathrm{Aut}(X^*(T))$ generated by $W(G,T)$ and $\phi_T(\mathrm{Gal}_K)$. So $\Pi(G,T)$ is isomorphic to a semidirect product $W(G,T) \rtimes \phi_T(\mathrm{Gal}_K)$.

In particular, if A is an abelian variety weakly of Mumford type of rank $2n+2$ and $\mathbf{G}_A$ the reductive group over $\mathbb{Q}$ constructed in [Pin98, Theorem 5.13], we have $W(\mathbf{G}_A, \mathbf{T}) \cong \{\pm 1\}^{2n+1}$. Let Ψ be a root system associated to $\mathbf{G}_A$. Since $\mathbf{G}_A^{\mathrm{der}}$ is semisimple, the group $\Pi(\mathbf{G}_A)$ is isomorphic to a subgroup of $\mathrm{Aut}(\Psi)$ which contains the Weyl group $W(\Psi) = W(\mathbf{G}_A) \cong \{\pm 1\}^{2n+1}$. The group $\mathrm{Aut}(\Psi)$ is isomorphic to the semidirect product $\{\pm 1\}^{2n+1} \rtimes S_{2n+1}$ where S_{2n+1} acts on $\{\pm 1\}^{2n+1}$ by permuting coordinates. So $\Pi(\mathbf{G}_A)$ is isomorphic to $\{\pm 1\}^{2n+1} \rtimes \phi_{\mathbf{T}}(\mathrm{Gal}_{\mathbb{Q}})$ where $\phi_{\mathbf{T}}(\mathrm{Gal}_{\mathbb{Q}})$ is some transitive subgroup of S_{2n+1}.

Let v be a place of good reduction and l any prime other than $p := \mathrm{char}(v)$. Since A has good reduction at v, there exists a polynomial $P_{A_v}(X) \in \mathbb{Q}[X]$ such that $P_{A_v}(X) = \det(XI - \rho_l(\mathrm{Fr}_v))$ for all $l \neq p$.

Now $A_{\overline{\mathbb{Q}}}$ is simple and the centralizer of $\mathbf{G}_A$ in $\mathrm{End}^0(\mathbf{V}_A)$ is isomorphic to $\mathrm{End}^0(A) = \mathbb{Q}$ by Faltings' theorem. Hence, the representation $\mathbf{G}_A \hookrightarrow \mathrm{GL}(\mathbf{V}_A)$ is irreducible. For each character

$\chi \in X^*(\mathbf{T})$, define $\mathbf{V}_A(\chi) = \{v \in \mathbf{V}_A \otimes_{\mathbb{Q}} \overline{\mathbb{Q}} : tv = \chi(t)v \ \forall\ t \in \mathbf{T}(\overline{\mathbb{Q}})\}$. Define the set of weights $\Omega = \{\chi \in X^*(\mathbf{T}) : \mathbf{V}_A(\chi) \neq 0\}$. We have the decomposition $\mathbf{V}_A \otimes_{\mathbb{Q}} \overline{\mathbb{Q}} = \bigoplus_{\chi \in \Omega} \mathbf{V}_A(\chi)$ and hence,

$$\det(XI - t) = \prod_{\chi \in \Omega} (X - \chi(t)).$$

The set Ω is stable under the actions of $W(\mathbf{G}_A, \mathbf{T})$ and $\mathrm{Gal}_{\mathbb{Q}}$ on $X^*(\mathbf{T})$ which gives a natural action of $\Pi(\mathbf{G}_A, \mathbf{T})$ on Ω. The representation $\mathbf{V}_A \otimes_{\mathbb{Q}} \overline{\mathbb{Q}}$ of $\mathbf{G}_{A,\overline{\mathbb{Q}}}$ is irreducible since $\mathrm{End}^0(A_{\overline{\mathbb{Q}}}) = \mathbb{Q}$ and $W(\mathbf{G}_A, \mathbf{T})$ acts transitively on Ω. The following result will be crucial for the density one result.

Theorem 5.8. *Let A be an abelian variety weakly of Mumford type of rank $2n+2$ over a number field F such that $F_A^{\mathrm{conn}} = F$. For any number field L splitting $\mathbf{G}_A$, we have $\mathrm{Gal}(L(\mathcal{W}_{A_v})/L) = W(\mathbf{G}_A, \mathbf{T})$ away from a set of places of density zero.*

The proof will cover the rest of this section. We will draw heavily from the techniques from [Zyw14] and [No09], although we will not assume the Mumford-Tate conjecture for our class of abelian varieties.

5.3 Frobenius conjugacy classes

The variety $\mathbf{Conj}'(\mathbf{G}_A)$. Let $G \subseteq \mathrm{GL}_{2g,\mathbb{Q}}$ be a connected reductive quasi-split algebraic group over $\mathbb{Q}$ such that its center is one-dimensional and G^{der} is $\mathbb{Q}$-simple. For simplicity, we assume that $\mathrm{Lie}(G^{\mathrm{der}})$ is $\otimes$-isotypic and the simple factors are of type A, B, C or $D^{\mathbb{H}}$. Let $G_1, \cdots, G_d$ be the simple factors of $G^{\mathrm{der}}_{\overline{\mathbb{Q}}}$. We have a central isogeny

$$G_1 \times \cdots \times G_d \longrightarrow G^{\mathrm{der}}_{\overline{\mathbb{Q}}}, \quad (g_1, g_2, \cdots, g_d) \mapsto g_1 g_2 \cdots g_d.$$

Let $\mathcal{A}$ be the group f of automorphisms of $G_{\overline{\mathbb{Q}}}$ such that
- f preserves each G_i.
- the restriction $f|_{G_i} : G_i \longrightarrow G_i$ agrees with conjugation by some element of G_i.
- f acts as the identity on the one-dimensional center of $G_{\overline{\mathbb{Q}}}$

Let R be the affine coordinate ring of G, on which G acts by composition with inner automorphisms. The group $\mathcal{A}$ acts on R by composition. We define $R^{\mathcal{A}}$ to be the $\mathbb{Q}$-subalgebra consisting of elements of elements of R fixed by $\mathcal{A}$. Now define $\mathbf{Conj}'(G) := \mathrm{Spec}(R^{\mathcal{A}})$. This is a universal categorical quotient. Let $\mathrm{Cl}'_G : G \longrightarrow \mathbf{Conj}'(G)$ be the morphism arising from the inclusion $R^{\mathcal{A}} \hookrightarrow R$ of $\mathbb{Q}$-algebras.

For a fixed maximal torus T of G, let $\mathcal{A}(T)$ be the subgroup of $\mathcal{A}$ that preserves $T_{\overline{\mathbb{Q}}}$. Define $\Gamma := \{f|_{T_{\overline{\mathbb{Q}}}} : f \in \mathcal{A}(T)\}$. This is a finite subgroup of $\mathrm{Aut}(T_{\overline{\mathbb{Q}}})$ stable under the $\mathrm{Gal}_{\mathbb{Q}}$-action. Furthermore, $\mathbf{Conj}'(G)$ is the quotient of $T_{\overline{\mathbb{Q}}}$ by Γ.

For a d-dimensional representation V of G, we have a morphism $P_V : \mathrm{GL}(V) \longrightarrow \mathbb{A}^d_{\mathbb{Q}}$ sending an element to the coefficients of its characteristic polynomial. This morphism factors through a map $\overline{P}_V : \mathbf{Conj}'(G) \longrightarrow \mathbb{A}^d_{\mathbb{Q}}$ which is a finite morphism of schemes.

Now let A be an abelian variety over a number field F, weakly of Mumford type of rank $2n+2$. Enlarging F if necessary, we may assume $F_A^{\mathrm{conn}} = F$. Since $\mathrm{End}(A_{\overline{\mathbb{Q}}}) = \mathbb{Z}$, there exists a quasi-split reductive group $\mathbf{G}_A$ over $\mathbb{Q}$ and a faithful $\mathbb{Q}$-linear representation $\mathbf{V}_A$ such that $\mathbf{G}_A^{\mathrm{der}}$ is $\mathbb{Q}$-simple and $(\mathbf{G}_A, \mathbf{V}_A) \otimes \mathbb{Q}_l \cong (G_l, V_l)$ for a set of rational primes of density one. The group $\mathbf{G}_A$ fulfills all of the conditions imposed on G. We define the variety $\mathbf{Conj}'(\mathbf{G}_A)$ as

described above. We also choose and fix a maximal torus $\mathbf{T}$ of $\mathbf{G}_A$. Note that in this case, $\Gamma = W(\mathbf{G}_A, \mathbf{T}) \cong \{\pm 1\}^{2n+1}$.

Definition 5.1. Following [No09], we say a place v of good reduction is *weakly neat* if none of the quotients of eigenvalues $\pi_1\pi_2^{-1}$ of the Frobenius are non-trivial roots of unity. They are *neat* if the group Φ_{A_v} is torsion-free.

The following definition is a straightforward generalization. Let G be a reductive group over a field K and V a faithful K-linear representation of G.

Definition 5.2. A semisimple element $g \in G(\overline{K})$ is *weakly neat* if none of the quotients of the eigenvalues of g are non-trivial roots of unity. It is *neat* if the group generated by the eigenvalues is torsion-free.

So a place v being weakly neat is equivalent to $\rho_l(\mathrm{Fr}_v)$ being weakly neat for any l other than $p = \mathrm{char}(v)$.

Remark Let B be an abelian variety B over a finite field k. Suppose $\mathcal{W}_B$ has two eigenvalues π_1, π_2 such that $(\pi_1\pi_2^{-1})^N = 1$. Then $\pi_1^N = \pi_2^N$. Let $\tilde{v}$ be a place lying over v in a finite extension over F such that the residue degree is N. Then $\mathcal{W}_{A_{\tilde{v}}} = \mathcal{W}_{A_v}^N$ has the eigenvalue π_1^N with multiplicity at least 2. Thus, for any abelian variety over a number field, all places of good ordinary reduction are weakly neat. In particular, for an abelian variety weakly of Mumford type, this condition holds away from a density zero set by [Pin98].

The following theorem is pivotal in the proof of Theorem 5.1. We recall (Theorem 2.4) that we have the inclusion $G_l^0(\mathbb{Q}_l) \subseteq \mathrm{MT}(\mathbb{Q}_l)$ for every prime l.

Theorem 5.9. ([No09], Theorem 1.8) *Let A be an abelian variety over a number field $K = K_A^{\mathrm{conn}}$. Let v be a weakly neat place of K. Then there exists an element $\mathrm{Cl}'_v \in \mathbf{Conj}'(\mathrm{MT}(A))(\mathbb{Q})$ such that $\mathrm{Cl}'_v = \mathrm{Cl}_{\mathrm{MT}(A)}(\rho_l(\mathrm{Fr}_v))$ for all primes l such that $v \nmid l$.*

Our approach will diverge here from that of [No09] and [Zyw14] since the Mumford-Tate conjecture remains open for abelian varieties weakly of Mumford type of rank greater than 4 (to our knowledge). We will need a variant of this compatibility theorem after replacing $\mathrm{MT}(A)$ by the group $\mathbf{G}_A$ and the set of all primes other than $\mathrm{char}(v)$ by the density one subset $\mathbb{S}_A$.

Let $\tau_n : \mathbf{G}_A \longrightarrow \mathbf{G}_A$ be the n-th power map. Since $\mathbf{Conj}'(\mathbf{G}_A)$ is a universal categorical quotient, there exists a unique map $\overline{\tau}_n : \mathbf{Conj}'(\mathbf{G}_A) \longrightarrow \mathbf{Conj}'(\mathbf{G}_A)$ making the following diagram commute.

$$\begin{array}{ccc} \mathbf{G}_A & \xrightarrow{\tau_n} & \mathbf{G}_A \\ \downarrow{\scriptstyle \mathrm{Cl}'_{\mathbf{G}_A}} & & \downarrow{\scriptstyle \mathrm{Cl}'_{\mathbf{G}_A}} \\ \mathbf{Conj}'(\mathbf{G}_A) & \xrightarrow{\overline{\tau}_n} & \mathbf{Conj}'(\mathbf{G}_A) \end{array}$$

We note that this is a finite map of $\mathbb{Q}$-varieties, which will be relevant later. For the next two lemmas, we drop the assumption that A is weakly of Mumford type and only assume the following:

1. $\mathrm{End}(A_{\overline{\mathbb{Q}}}) = \mathbb{Z}$. So the triple $(\mathbf{G}_A, \mathbf{V}_A, \mathbb{S}_A)$ is well-defined and $\mathbf{G}_A^{\mathrm{der}}$ is $\mathbb{Q}$-simple.
2. The root systems of $\mathrm{Lie}(\mathbf{G}_{A,\overline{\mathbb{Q}}}^{\mathrm{der}})$ are of type A, B, C or $D^{\mathbb{H}}$.

Alternatively, we are assuming that the root systems of $\mathrm{Lie}(\mathbf{G}_{A,\overline{\mathbb{Q}}}^{\mathrm{der}})$ are not of type $D^{\mathbb{R}}$.

Lemma 5.10. *Let α, β be two weakly neat elements of $\mathbf{G}_A(\overline{\mathbb{Q}})$ with the same characteristic polynomial in $\mathbf{V}_A$. If $\overline{\tau}_n(\alpha) = \overline{\tau}_n(\beta)$, then $\mathrm{Cl}'_{\mathbf{G}_A}(\alpha) = \mathrm{Cl}'_{\mathbf{G}_A}(\beta)$.*

Proof. Write $g = \dim A$ for brevity. The representation $\mathbf{V}_A$ is $2g$-dimensional and the symmetric group S_{2g} acts on $\mathbb{G}^{2g}_{m,\overline{\mathbb{Q}}}$ by permutation. Define the set

$$\mathbf{T}_w := \{t \in \mathbf{T} : t \text{ is weakly neat and } \mathrm{Cent}_{S_{2g}}(t) = \mathrm{Cent}_{S_{2g}}(\mathbf{T})\}.$$

Let $s, t \in \mathbf{T}_w$ be elements such that $\tau_n(s) = \tau_n(t)$ and s, t have the same characteristic polynomial. Then there exists some $\tau \in S_g$ such that $\tau(s) = t$. Thus, $\tau(s^n) = t^n$ and hence $\tau \in \mathrm{Cent}_{S_{2g}}(\mathbf{T})$. So τ is the identity on $\mathbf{T}$ and hence, $t_1 = t_2$.

Since α is a semisimple element, it lies in a maximal torus in $\mathrm{GL}(\mathbf{V}_A)$. Write $\alpha = (\alpha_1, \cdots, \alpha_{2g}) \in \mathbb{G}^{2g}_m$. We define the torus $\mathbf{T}^\alpha = \{(t_1, \cdots, t_{2g}) \in \mathbb{G}^{2g}_m : t_i = t_j \text{ if } \alpha_i = \alpha_j\}$. Then we have $\alpha \in \mathbf{T}^\alpha_w$. Since $\mathrm{Cl}'_{\mathbf{G}_A}(\alpha^n) = \mathrm{Cl}'_{\mathbf{G}_A}(\beta^n)$, it follows that α^n and β^n are conjugate in $\mathbf{G}_A(\overline{\mathbb{Q}})$. So there exists a weakly neat semisimple element $\gamma \in \mathbf{G}_A(\overline{\mathbb{Q}})$ such that $\alpha^n = \gamma^n$ and $\mathrm{Cl}'_{\mathbf{G}_A}(\beta) = \mathrm{Cl}'_{\mathbf{G}_A}(\gamma)$. So α and γ have the same characteristic polynomial. Since $\alpha^n = \gamma^n$, we have $\gamma \in T^\alpha_w$. Thus, $\mathrm{Cl}'_{\mathbf{G}_A}(\alpha) = \mathrm{Cl}'_{\mathbf{G}_A}(\beta)$. □

In particular, the map

$$\tau_n \times P_V : \mathbf{Conj}'(\mathbf{G}_A) \longrightarrow \mathbf{Conj}'(\mathbf{G}_A) \times \mathbb{A}^{2g}_{\mathbb{Q}}$$

is injective on the subset of $\mathbf{Conj}'(\mathbf{G}_A)(\overline{\mathbb{Q}})$ that is the image of the subset of weakly neat elements in $\mathbf{G}_A(\overline{\mathbb{Q}})$. We now use this lemma to show that the theorem is true if it is "potentially" true, in the sense of passing to a finite extension.

Proposition 5.11. *Let A be an abelian variety over a number field F such that $F_A^{\mathrm{conn}} = F$ and $\mathrm{End}(A_{\overline{\mathbb{Q}}}) = \mathbb{Z}$. Let v be a weakly neat place of F. Suppose there exists a finite extension F'/F with a place $\tilde{v}$ lying over v such that for some conjugacy class $\mathrm{Cl}'_{\tilde{v}} \in \mathbf{Conj}'(\mathbf{G}_A)(\mathbb{Q})$, we have $\mathrm{Cl}'_{\mathbf{G}_A}(\rho_l(\mathrm{Fr}_{\tilde{v}})) = \mathrm{Cl}'_{\tilde{v}}$ for all $l \in \mathbb{S}_A$. Then there exists a conjugacy class $\mathrm{Cl}'_v \in \mathbf{Conj}'(\mathbf{G}_A)(\mathbb{Q})$ such that $\mathrm{Cl}'_{\mathbf{G}_A}(\rho_l(\mathrm{Fr}_v)) = \mathrm{Cl}'_v$ for all $l \in \mathbb{S}_A$.*

Proof. Let $l \in \mathbb{S}_A$. Pick an element $x := \mathrm{Cl}'_{\mathbf{G}_A}(\mathrm{Fr}_v) \in \mathbf{Conj}'\mathbf{G}_A(\mathbb{Q}) \subseteq \mathbf{Conj}'\mathbf{G}_A(\overline{\mathbb{Q}}_l)$. Let n be the residue degree of $\tilde{v}$ over v. Let $\tilde{x}$ be a pre-image of $\mathrm{Cl}'_{\mathbf{G}_A}(\mathrm{Fr}_v)$ under the map of $\overline{\mathbb{Q}}_l$-valued points $\mathbf{Conj}'(\mathbf{G}_A)(\overline{\mathbb{Q}}_l) \longrightarrow \mathrm{Cl}'_{\mathbf{G}_A}(\overline{\mathbb{Q}}_l)$ induced by $\overline{\tau}_n$. We first show that $\tilde{x}$ is a $\overline{\mathbb{Q}}$-valued point of $\mathbf{Conj}'(\mathbf{G}_A)(\overline{\mathbb{Q}}_l)$.

Let $R^{\mathcal{A}}$ be the coordinate ring of $\mathbf{Conj}'(\mathbf{G}_A)$ and let $\overline{\tau}_n^\sharp : R^{\mathcal{A}} \longrightarrow R^{\mathcal{A}}$ be the homomorphism induced on the rings. Then $\tau_n \circ \tilde{x} = x$. So we have the commutative diagram:

$$\begin{array}{ccc} R^{\mathcal{A}} & \xrightarrow{\overline{\tau}_n^\sharp} & R^{\mathcal{A}} \\ \downarrow x & & \downarrow \tilde{x} \\ \mathbb{Q} & \xrightarrow{i} & \overline{\mathbb{Q}}_l \end{array}$$

Since $\tau_n^\sharp$ is a finite homomorphism of algebras, $\tilde{x}(R^{\mathcal{A}})$ is a finite algebra over $x(R^{\mathcal{A}}) = \mathbb{Q}$. So $K := \mathrm{Frac}(\tilde{x}(R^{\mathcal{A}}))$ is a number field. The diagram above may be replaced by:

$$\begin{array}{ccc} R^{\mathcal{A}} & \xrightarrow{\overline{\tau}_n^\sharp} & R^{\mathcal{A}} \\ \downarrow x & & \downarrow \tilde{x} \\ \mathbb{Q} & \xrightarrow{i} & K \end{array}$$

Since $\tilde{x}$ is a K-valued point of $\mathbf{Conj}'(\mathbf{G}_A)$, this proves the claim that $\tilde{x}$ is a $\overline{\mathbb{Q}}$-valued point of $\mathbf{Conj}'(\mathbf{G}_A)$.

Now $\mathrm{Cl}'_{\mathbf{G}_A}(\rho_{v,l}(\mathrm{Fr}_v))$ is a $\mathbb{Q}_l$-valued point of $\mathbf{Conj}'(\mathbf{G}_A)$ for every $l(\neq p)$ in $\mathbb{S}_A$. Let l, l' be distinct primes in $\mathbb{S}_A$ other than p. Then we have a weakly neat element $\rho_{l'}(\mathrm{Fr}_v)$ with the same characteristic polynomial as that of $\rho_l(\mathrm{Fr}_v)$. By the hypothesis, we have an element $\mathrm{Cl}'_{\mathbf{G}_A}(\rho_l(\mathrm{Fr}_v)) \in \mathbf{Conj}'(\mathbf{G}_A)(\mathbb{Q}_l)$ such that

$$\overline{\tau}_n(\mathrm{Cl}'_{\mathbf{G}_A}(\rho_l(\mathrm{Fr}_v))) = \overline{\tau}_n(\mathrm{Cl}'_{\mathbf{G}_A}(\rho_{l'}(\mathrm{Fr}_v))).$$

By the preceding discussion, $\mathrm{Cl}'_{\mathbf{G}_A}(\rho_{l'}(\mathrm{Fr}_v)) \in \mathbf{Conj}'(\mathbf{G}_A)(\overline{\mathbb{Q}})$. So by lemma 5.10, it follows that $\mathrm{Cl}'_{\mathbf{G}_A}(\rho_l(\mathrm{Fr}_v)) = \mathrm{Cl}'_{\mathbf{G}_A}(\rho_{l'}(\mathrm{Fr}_v))$ for all $l, l' \in \mathbb{S}_A$ other than p. We denote this element by Cl'_v.

It remains to show that Cl'_v is $\mathbb{Q}$-valued. Now, $\mathrm{Cl}'_v \in \mathbf{Conj}'(\mathbf{G}_A)(\mathbb{Q}_l)$ for all $l \in \mathbb{S}_A$ other than p. Suppose, by way of contradiction, that $\mathrm{Cl}'_v \notin \mathbf{Conj}'(\mathbf{G}_A)(\mathbb{Q})$ and let $K \neq \mathbb{Q}$ be the smallest number field such that $\mathrm{Cl}'_v \in \mathbf{Conj}'(\mathbf{G}_A)(K)$. Write $K = \mathbb{Q}(\alpha)$ with $\alpha \in \mathcal{O}_K$ and let f be the irreducible polynomial of α over $\mathbb{Q}$ with $\deg(f) \geq 2$. By the Chebotarev density theorem, the set of primes

$$\mathcal{P} := \{l : \text{ The image of } f(X) \text{ in } \mathbb{F}_l[X] \text{ is separable and has no zero}\}$$

has positive density. For any $l \in \mathcal{P}$, Hensel's lemma implies that K does not have an embedding in $\mathbb{Q}_l$. Since $\mathbb{S}_A$ is of Dirichlet density one and is disjoint from $\mathcal{P}$, we have a contradiction. Thus, $\mathrm{Cl}'_v \in \mathbf{Conj}'(\mathbf{G}_A)(\mathbb{Q})$. □

Proposition 5.12. *Let A be an abelian variety weakly of Mumford type of rank $2n+2$ over a number field $F = F_A^{\mathrm{conn}}$. Let $(\mathbf{G}_A, \mathbf{V}_A, \mathbb{S}_A)$ be a triple as in* ([Pin98], Theorem 5.1)*. Let v be a weakly neat place of F. Then there exists a conjugacy class $\mathrm{Cl}'_v \in \mathbf{Conj}'(\mathbf{G}_A)(\mathbb{Q})$ such that $\mathrm{Cl}'_{\mathbf{G}_A}(\rho_l(\mathrm{Fr}_v)) = \mathrm{Cl}'_v$ for all $l \in \mathbb{S}_A$ other than $p = \mathrm{char}(v)$.*

Proof. Since $\mathrm{End}(A_{\overline{\mathbb{Q}}}) = \mathbb{Z}$, the $\mathbf{G}_{A,\overline{\mathbb{Q}}}$-representation $\mathbf{V}_A \otimes_{\mathbb{Q}} \overline{\mathbb{Q}}$ is irreducible. The group of homotheties of $\mathbf{V}_A \otimes_{\mathbb{Q}} \overline{\mathbb{Q}}$ commutes with the action of $\mathbf{G}^{\mathrm{der}}_{A,\overline{\mathbb{Q}}}$ which is isomorphic to $\mathrm{SL}^{2n+1}_{2,\overline{\mathbb{Q}}}$. We write $g = 2^{2n}$. Define the map $P_{\mathbf{V}_A} : \mathbf{G}_A(\overline{\mathbb{Q}}) \longrightarrow \mathbb{A}^{2g}_{\mathbb{Q}}(\overline{\mathbb{Q}})$ sending an element to the coefficients of its characteristic polynomial. It factors through a finite morphism $\overline{P}_{\mathbf{V}_A} : \mathbf{Conj}'(\mathbf{G}_A) \longrightarrow \mathbb{A}^{2g}_{\mathbb{Q}}$. Let $\tilde{S}_{w.n.} \subseteq \mathbf{G}_A(\overline{\mathbb{Q}})$ be the subset of weakly neat elements and let $S_{w.n.}$ be its image in $\mathbf{Conj}'(\mathbf{G}_A)(\overline{\mathbb{Q}})$ under the map $\mathrm{Cl}'_{\mathbf{G}_A}$.

Now $\mathbf{G}^{\mathrm{der}}_{A,\overline{\mathbb{Q}}} \cong \mathrm{SL}^{2n+1}_{2,\overline{\mathbb{Q}}}$ and in particular, $\mathbf{G}^{\mathrm{der}}_{A,\overline{\mathbb{Q}}}$ is $\otimes$-isotypic of type A. So the characteristic polynomial determines the conjugacy class of a semisimple element. In particular, the map $\overline{P}_{\mathbf{V}_A} : \mathbf{Conj}'(\mathbf{G}_A) \longrightarrow \mathbb{A}^{2g}_{\mathbb{Q}}$ is injective on the subset $S_{w.n.}$. Hence, there is a unique element $\mathrm{Cl}'_v \in \mathbf{Conj}'(\mathbf{G}_A)(\mathbb{Q})$ such that $\overline{P}_{\mathbf{V}_A}(\mathrm{Cl}'_v) = P_{\mathbf{V}_A}(\mathrm{Fr}_v)$. Thus, $\mathrm{Cl}'_v = \mathrm{Cl}'_{\mathbf{G}_A}(\rho_l(\mathrm{Fr}_v))$ for all $l \in \mathbb{S}_A$. Furthermore, since the common characteristic polynomials of all the elements $\rho_l(\mathrm{Fr}_v)$, $v \nmid l$ has rational coefficients, it follows that $\mathrm{Cl}'_v \in \mathbf{Conj}'(\mathbf{G}_A)(\mathbb{Q})$. □

5.4 Galois representations

For a place v of F, we have the surjective homomorphism $\mathrm{ev} : X^*(\mathbf{T}) \longrightarrow \Phi_{A_v}$, $\chi \mapsto \chi(t_v)$ that maps Ω onto $\mathcal{W}_{A_v}$. For places in $\mathcal{S}_A$, it is an isomorphism.

Lemma 5.13. *Let $v \in \mathcal{S}_A$ and fix an element $t \in \mathbf{T}(\overline{\mathbb{Q}})$ such that $\det(XI - t) = P_{A_v}(X)$. Then the map*

$$\mathrm{ev} : X^*(\mathbf{T}) \longrightarrow \Phi_{A_v}, \quad \chi \mapsto \chi(t)$$

is an isomorphism mapping Ω onto $\mathcal{W}_{A_v}$.

Proof. It is clear that the map is a homomorphism $X^*(\mathbf{T}) \longrightarrow \overline{\mathbb{Q}}^*$. The zeros of $P_{A_v}(X)$ are the values $\chi(t)$ with $\chi \in \Omega$. So $\mathcal{W}_{A_v} = \mathrm{ev}(\Omega)$. The set Ω generates $X^*(\mathbf{T})$ since $\mathbf{G}_A$ acts faithfully on $\mathbf{V}_A$. Since $\mathrm{ev}(\Omega) = \mathcal{W}_{A_v}$ and $\mathcal{W}_{A_v}$ generates Φ_{A_v}, it follows that $\mathrm{ev}(X^*(\mathbf{T})) = \Phi_{A_v}$. Thus, ev is a surjective homomorphism of free abelian groups. So it suffices to verify that the ranks of both groups coincide.

Since $v \in \mathcal{S}_A$, rank of the group Φ_{A_v} is $\mathrm{rank}(A)$. On the other hand, $X^*(\mathbf{T})$ is a free abelian group of the same rank as $\mathbf{G}_A^{\mathrm{der}}$, which is $2n+1$. Since ev is surjective, it follows that it is an isomorphism. □

The upshot is that for any $v \in \mathcal{S}_A$, if $W(\mathbf{G}_A, \mathbf{T}) \subseteq \mathrm{Gal}(\mathbb{Q}(\mathcal{W}_{A_v})/\mathbb{Q})$, then $\mathrm{Gal}_{\mathbb{Q}}$ acts transitively on $\mathcal{W}_{A_v}$ and hence, the reduction is absolutely simple. Thus, detecting the places of simple reduction boils down to detecting places where $W(\mathbf{G}_A, \mathbf{T}) \subseteq \mathrm{Gal}(\mathbb{Q}(\mathcal{W}_{A_v})/\mathbb{Q})$.

Fix a place $v \in \mathcal{S}_A$. By Proposition 5.12, we may choose an element $t_v \in \mathbf{T}(\overline{\mathbb{Q}})$ such that $\mathrm{Cl}'_{\mathbf{G}_A}(t_v) = \mathrm{Cl}'_v \in \mathbf{Conj}'(\mathbf{G}_A)(\mathbb{Q})$. We may choose t_v so that $\det(XI - t_v) = P_{A_v}(X)$. We define $\psi_v(\sigma)$ to be the unique element in $\mathrm{Aut}(X^*(\mathbf{T}))$ such that the following diagram commutes:

$$\begin{array}{ccc} X^*(\mathbf{T}) & \xrightarrow{\mathrm{ev}} & \Phi_{A_v} \\ \big\downarrow{\scriptstyle \psi_v(\sigma)} & & \big\downarrow{\scriptstyle \sigma} \\ X^*(\mathbf{T}) & \xrightarrow{\mathrm{ev}} & \Phi_{A_v} \end{array}$$

This yields the Galois representation $\psi_v : \mathrm{Gal}_{\mathbb{Q}} \longrightarrow \mathrm{Aut}(X^*(\mathbf{T}))$, $\sigma \mapsto \psi_v(\sigma)$. Now $W(\mathbf{G}_A, \mathbf{T}) \cong W(G_l, \mathbf{T}_{\mathbb{Q}_l})$. If $\mathbf{T}_{\mathbb{Q}_l}$ is split and $v \nmid l$, we have a group homomorphism $\psi_{v,l} : \mathrm{Gal}_{\mathbb{Q}_l} \longrightarrow W(G_l, \mathbf{T}_l)$ such that $\sigma(t_{v,l}) = \psi_{v,l}(\sigma)^{-1}(t_{v,l})$. This map is independent of the choice of $t_{v,l}$.

Remark. Let K be a number field splitting $\mathbf{T}$. Note that if l is split completely in K, then $K_w = \mathbb{Q}_l$ for any prime w of $\mathbb{Q}(\mathcal{W}_{A_v})$ lying over l. Hence, $\mathbf{T}_{\mathbb{Q}_l}$ is split. Since $\mathbb{S}_A$ has density one, the set of primes $l \in \mathbb{S}_A$ such that $\mathbf{T}_{\mathbb{Q}_l}$ is split has positive density.

Lemma 5.14. *Let $l \in \mathbb{S}_A$ and suppose $\mathbf{T}_{\mathbb{Q}_l}$ is split. For any $\sigma \in \mathrm{Gal}_{\mathbb{Q}_l}$, we have $\psi_v(\sigma) = \psi_{v,l}(\sigma) \in W(\mathbf{G}_A, \mathbf{T})$*

Proof. Choose an element $t_{v,l} \in \mathbf{T}(\overline{\mathbb{Q}}_l)$ conjugate to $\rho_l(\mathrm{Fr}_v)$ in $\mathbf{G}_A(\overline{\mathbb{Q}}_l)$. Then $\mathrm{Cl}'_{\mathbf{G}_A}(t_{v,l}) = \mathrm{Cl}'_{\mathbf{G}_A}(\rho_l(\mathrm{Fr}_v)) = \mathrm{Cl}'_v$. So there is a unique $\beta \in \Gamma = W(\mathbf{G}_A, \mathbf{T})$ such that $t_{v,l} = \beta(t_v)$. We may check that $\psi_v(\sigma) = \beta^{-1}\psi_{v,l}(\sigma) \circ \beta$. Since $\psi_{v,l}(\sigma) \in W(\mathbf{G}_A, \mathbf{T})$ and $W(\mathbf{G}_A, \mathbf{T})$ is abelian in this particular case, we have $\psi_v(\sigma) = \psi_{v,l}(\sigma) \in W(\mathbf{G}_A, \mathbf{T})$. □

Let l be a prime such that $\mathcal{G}_l$ is a reductive group scheme over $\mathbb{Z}_l$, as is the case when l is sufficiently large. Choose a maximal torus $\mathcal{T}_l$ in $\mathcal{G}_l$ and let $\mathbf{T}_l$ be its generic fiber, which is a maximal torus in G_l. For any place $v \in \mathcal{S}_A$ such that $v \nmid l$, we define the conjugacy class

$$\mathcal{I}_{v,l} := \{t \in \mathbf{T}(\overline{\mathbb{Q}}_l) : t \text{ and } \rho_l(\mathrm{Fr}_v) \text{ are conjugate in } G_l(\overline{\mathbb{Q}}_l)\}$$

and fix an element $t_{v,l} \in \mathcal{I}_{v,l}$. The third condition in the definition of the set $\mathcal{S}_A$ ensures that the group generated by $t_{v,l}$ is Zariski dense in $\mathbf{T}_{l,\overline{\mathbb{Q}}_l}$. So the action of $W(G_l, \mathbf{T}_l)$ on $\mathcal{I}_{v,l}$ is simply transitive.

For each $\sigma \in \mathrm{Gal}_{\mathbb{Q}_l}$, there is a unique $\psi_{v,l}(\sigma) \in W(G_l, \mathbf{T}_l)$ such that $\sigma(t_{v,l}) = \psi_{v,l}(\sigma)^{-1}(t_{v,l})$. Since $\mathbf{T}_l$ is split by assumption, the map

$$\psi_{v,l} : \mathrm{Gal}_{\mathbb{Q}_l} \longrightarrow W(G_l, \mathbf{T}_l), \quad \sigma \mapsto \psi_{v,l}(\sigma)$$

is a group homomorphism.

Notation: Define the field $k_{\mathbf{G}_A}$ to be the intersection of all algebraic extensions of $\mathbb{Q}$ that split $\mathbf{G}_A$. Since $\mathbf{G}_A$ is quasi-split ([Pin98], Theorem 5.13), the field $k_{\mathbf{G}_A}$ is a number field. We first verify the inclusion $\psi_v(\mathrm{Gal}_{k_{\mathbf{G}_A}}) \subseteq W(\mathbf{G}_A, \mathbf{T})$.

Lemma 5.15. *The image $\psi_v(\mathrm{Gal}_{k_{\mathbf{G}_A}})$ is a subgroup of $W(\mathbf{G}_A, \mathbf{T})$.*

Proof. It suffices to show that $\psi_v(\mathrm{Gal}_L) \subseteq W(\mathbf{G}_A, \mathbf{T})$ for any number field L splitting $\mathbf{G}_A$.

Let L be a number field that splits $\mathbf{T}$. Let Λ be the co-finite subset of $\mathcal{S}(L|\mathbb{Q}) \cap \mathbb{S}_A$ of primes l for which ψ_v is unramified at l, $v \nmid l$. The torus $\mathbf{T}_{\mathbb{Q}_l}$ is split for all $l \in \Lambda$. From the preceding lemma, $\psi_v(\mathrm{Fr}_l) \in W(\mathbf{G}_A, \mathbf{T})$. Since, the set $\{\mathrm{Fr}_l : l \in \Lambda\}$ generates Gal_L, it follows that $\psi_v(\mathrm{Gal}_L) \subseteq W(\mathbf{G}_A, \mathbf{T})$.

Now let L be a number field splitting $\mathbf{G}_A$. Choose a maximal torus $\mathbf{T}'$ of $\mathbf{G}_A$ for which $\mathbf{T}'_L$ is split and an element $g \in \mathbf{G}_A(\overline{\mathbb{Q}})$ such that $\mathbf{T}'_{\overline{\mathbb{Q}}} = g\mathbf{T}_{\overline{\mathbb{Q}}}g^{-1}$. Define $t'_v := gt_vg^{-1}$. We have $\mathrm{Cl}'_{\mathbf{G}_A}(t'_v) = \mathrm{Cl}'_v$ and $\det(XI - t'_v) = P_{A_v}(X)$. We define the homomorphism $\psi'_v : \mathrm{Gal}_{\mathbb{Q}} \longrightarrow \mathrm{Aut}(X^*(\mathbf{T}'))$ such that $\sigma(\alpha(t'_v)) = (\psi'_v\alpha)(t'_v)$ for all $\alpha \in X^*(\mathbf{T})$ and $\sigma \in \mathrm{Gal}_{\mathbb{Q}}$.

Let $\gamma : \mathbf{T}_{\overline{\mathbb{Q}}} \longrightarrow \mathbf{T}'_{\overline{\mathbb{Q}}}$ be the conjugation by g and let $\gamma_* : \mathrm{Aut}(\mathbf{T}_{\overline{\mathbb{Q}}}) \longrightarrow \mathrm{Aut}(\mathbf{T}'_{\overline{\mathbb{Q}}})$ be the conjugation by γ. Then $\gamma_*(W(\mathbf{G}_A, \mathbf{T})) = W(\mathbf{G}_A, \mathbf{T}')$. From the definition of $\psi_v(\sigma)$, we check that $\psi_v(\sigma) = \gamma^{-1} \circ \psi'_v(\sigma) \circ \gamma = \gamma_*^{-1}(\psi'_v(\sigma))$. So $\psi_v(\mathrm{Gal}_L) \subseteq \gamma_*^{-1}(W(\mathbf{G}_A, \mathbf{T}')) = W(\mathbf{G}_A, \mathbf{T})$. □

The image of ψ_v is, in fact, the full $W(\mathbf{G}_A, \mathbf{T})$ for almost all places v, which will complete the proof of the main result. The following lemma from [Zyw14] will be necessary.

Lemma 5.16. ([Zyw14], Lemma 5.1) *Fix a subset C of $W(G_l, \mathbf{T}_{\mathbb{Q}_l})$ stable under conjugation. There is a subset U_l of $\rho_l(\mathrm{Gal}_F)$ stable under conjugation with the following properties:*

(a). If $v \in \mathcal{S}_A$ satisfies $v \nmid l$ and $\overline{\rho}_l(\mathrm{Fr}_v) \subseteq U_l$, then $\psi_{v,l}$ is unramified and $\psi_{v,l}(\mathrm{Fr}_l) \subseteq C$.
(b). Let F' be a finite extension of F and let H be a coset of $\mathrm{Gal}_{F'}$ in Gal_F. Then we have

$$\frac{\big|\overline{\rho}_l(H) \cap U_l\big|}{\big|\overline{\rho}_l(H)\big|} = \frac{\big|C\big|}{W(G_l, \mathbf{T}_{\mathbb{Q}_l})} + O(1/l)$$

where the implicit constant depends only on A and F'.

The main ideas in the following proposition are from ([Zyw14], Proposition 6.6). To avoid confusion with the notation, we reiterate that we use the symbol $\mathbf{G}_A$ for the group constructed in ([Pin98], Theorem 5.13) rather than the Mumford-Tate group $\mathrm{MT}(A)$.

Proposition 5.17. *Let A be an abelian variety weakly of Mumford type over a number field F such that $F = F_A^{\mathrm{conn}}$. Let L be a finite extension of $k_{\mathbf{G}_A}$. Then $\psi_v(\mathrm{Gal}_L) = W(\mathbf{G}_A, \mathbf{T})$ for all places $v \in \mathcal{S}_A$ away from a subset of Dirichlet density 0.*

Proof. Fix a full rank $\mathbf{G}_A$-stable lattice $\mathcal{L} \subseteq \mathbf{V}_A$. Let $\mathcal{T}$ be the Zariski closure of $\mathbf{T}$ in the group scheme $\mathrm{GL}(\mathcal{L})$. For l sufficiently large, $\mathcal{T}_{\mathbb{Z}_l}$ is a torus over $\mathbb{Z}_l$, by [LP95]. Since rank $\mathbf{G}_A$ = rank G_l, it follows that $\mathcal{T}_{\mathbb{Z}_l}$ is, in fact, a maximal torus of $\mathcal{G}_l$ for l sufficiently large and in $\mathbb{S}_A$.

Let F'/F be a finite extension fulfilling the conditions of Serre's lemma. Let c be an element of $W(\mathbf{G}_A, \mathbf{T})$. Since $\Gamma = W(\mathbf{G}_A, \mathbf{T}) \cong \{\pm 1\}^{2n+1}$, the singleton set $\{c\}$ is stable under conjugation

by Γ in this case. Let $\mathcal{S}(L|\mathbb{Q})$ be the set of rational primes split completely in L. For l in the set $\mathcal{S}(L|\mathbb{Q})\cap\mathbb{S}_A$ and sufficiently large, we may identify c with an element of $W(G_l, \mathbf{T}_{\mathbb{Q}_l}) = W(\mathcal{G}_l, \mathcal{T}_{\mathbb{Q}_l})$. Let $\mathcal{V}$ be the subset of $\mathcal{S}_A$ of places v such that $\overline{\rho}_l(\mathrm{Fr}_v) \nsubseteq U_l$, $l \neq p$. Let $\mathcal{V}_Q$ the set of places such that $\overline{\rho}_l(\mathrm{Fr}_v) \nsubseteq U_l$ for all $l < Q$, $l \neq p$. Then $\mathcal{V} = \bigcap_{Q=1}^{\infty} \mathcal{V}_Q$. By ([Zyw14], Proposition 2.12), the set $\mathcal{V}_Q$ has density

$$\sum_{\mathcal{C}} \frac{|\mathcal{C}|}{\mathrm{Gal}(F'/F)} \prod_{l\in\mathcal{S}(L|\mathbb{Q})\,\cap\,\mathbb{S}_A\,\cap[0,Q]} \frac{\left|\overline{\rho}_l(\Gamma_{\mathcal{C}}) \cap (\rho_l(\mathrm{Gal}_F) \setminus U_l)\right|}{\left|\overline{\rho}_l(\Gamma_{\mathcal{C}})\right|}$$

where $\mathcal{C}$ varies over the conjugacy classes of $\mathrm{Gal}(F'/F)$ and $\Gamma_{\mathcal{C}} = \{\sigma \in \mathrm{Gal}_F : \sigma|_{F'} \in \mathcal{C}\}$. Furthermore, by ([Zyw14], Lemma 5.1 (b)),

$$\begin{aligned}\mathrm{Density}(\mathcal{V}_Q) &<< \prod_{l\in\mathcal{S}(L|\mathbb{Q})\,\cap\,\mathbb{S}_A\,\cap\,[0,Q]} \left(1 - \frac{1}{W(G_l, \mathbf{T}_{\mathbb{Q}_l})} + O(1/l)\right) \\ &= \prod_{l\in\mathcal{S}(L|\mathbb{Q})\,\cap\,\mathbb{S}_A\cap\,[0,Q]} \left(1 - \frac{1}{W(\mathbf{G}_A, \mathbf{T})} + O(1/l)\right).\end{aligned}$$

Now, $\mathcal{S}(L|\mathbb{Q}) \cap \mathbb{S}_A$ has density $\frac{1}{[\tilde{L}:\mathbb{Q}]}$ where $\tilde{L}$ is the Galois closure of L over $\mathbb{Q}$. In particular, the set $\mathcal{S}(L|\mathbb{Q}) \cap \mathbb{S}_A$ is infinite and hence, $\mathcal{V}$ has density zero. Let $v \in \mathcal{S}_A \setminus \mathcal{V}$. Choose a prime $l \in \mathcal{S}(L|\mathbb{Q}) \cap \mathbb{S}_A$ such that $v \nmid l$ and $\overline{\rho}_l(\mathrm{Fr}_v) \subseteq U_l$. Then $\psi_{v,l}$ is unramified at l and $\psi_{v,l}(\mathrm{Fr}_v) = c$ by ([Zyw14], Lemma 5.1(a)). Since $\{c\}$ is stable under conjugation by Γ and the elements $\psi_v(\mathrm{Fr}_l) = \psi_{v,l}(\mathrm{Fr}_l) \in W(\mathbf{G}_A, \mathbf{T})$, it follows that $\psi_v(\mathrm{Fr}_l) = c$. Since c was an arbitrary element of $W(\mathbf{G}_A, \mathbf{T})$, we have $\psi_v(\mathrm{Gal}_L) = W(\mathbf{G}_A, \mathbf{T})$. □

The main result (Theorem 1.1) now follows:

Proof. Let L be any number field splitting $\mathbf{G}_A$. Suppose v lies in the density one subset of $\mathcal{S}_A$ described in the preceding Proposition. Since $\mathcal{W}_{A_v}$ generates Φ_{A_v}, the restriction $\psi_v|_{\mathrm{Gal}_L}$ factors through an isomorphism $\mathrm{Gal}(L(\mathcal{W}_{A_v})/L) \xrightarrow{\cong} W(\mathbf{G}_A)$. So $\mathrm{Gal}_{\mathbb{Q}}$ acts transitively on $\mathcal{W}_{A_v}$ and hence, A_v is absolutely simple. □

6 Appendix

6.1 Generalized Mumford Curves

We verify that abelian varieties with Mumford-Tate groups fulfilling the hypothesis of Definition 1.3 arise as generic points on Mumford curves. The proof is identical to that of [No01, Proposition 1.5] for the case $n = 1$. We provide a sketch for the reader's convenience.

Proposition 6.1. *Let X be a an abelian variety of dimension 2^{2n}, $V = H^1(X(\mathbb{C}), \mathbb{Q})$ the Betti cohomology and $G = \mathrm{MT}(X)$ its Mumford-Tate group. Assume (G, V) is of Mumford type of rank $2n + 2$. Then there exist a totally real number field K of degree $2n + 1$ and a quaternion D/K such that:*
(i) *D is split at exactly one real place*
(ii) $\mathrm{Cor}_{K/\mathbb{Q}}(D) = 0$
(iii) *X is isomorphic to a fiber of the family $\mathcal{X}/M$ over a $\mathbb{C}$-valued point of $M = M_C$ (for any sufficiently small C).*

Proof. Let $\tilde{G} \longrightarrow G$ be the universal simple cover with $\tilde{G} \cong \mathbb{G}_{m,\mathbb{Q}} \times \tilde{G}'$, $\tilde{G}'_{\overline{\mathbb{Q}}} \cong \mathrm{SL}_{2,\overline{\mathbb{Q}}}^{2n+1}$. Now G^{der} is $\mathbb{Q}$-simple. So $\mathrm{Gal}_{\mathbb{Q}}$ acts transitively on the set of factors of the product $\mathrm{SL}_{2,\overline{\mathbb{Q}}}^{2n+1}$. Let $H \subseteq \mathrm{Gal}_{\mathbb{Q}}$ be the subgroup fixing the first factor and let $K := \overline{\mathbb{Q}}^H$. Then K is field of degree $2n+1$ and G' is the Weil restriction from K to $\mathbb{Q}$ of a K-form of SL_2. By Kneser's theorem ([Kne65], Satz 3), there exists a central division algebra D of degree 4 over K such that $\tilde{G}' = (D^*)^{\mathrm{der}}$. Since $\tilde{G}'$ acts on a 2^{2n+1}-dimensional $\mathbb{Q}$-vector space, it follows that $\mathrm{Cor}_{K/\mathbb{Q}}(D) = \mathrm{Mat}_{2^{2n+1}}(\mathbb{Q})$.

Now V carries a symplectic form $\langle\ .\ ,.\ \rangle$. Let $h : S \longrightarrow G_{\mathbb{R}}$ be the morphism defining the Hodge structure on $V \otimes \mathbb{R}$. The symmetric bilinear form $\langle\ .\ , h(i).\ \rangle$ is positive definite. This implies that if $\mathcal{H}$ is the real form of $G_{\mathbb{C}}$ corresponding to the involution $\mathrm{ad}(h(i))$, then $\mathcal{H}^{\mathrm{der}}$ is compact.

The projection of h on precisely one factor of $\mathrm{PSL}_{2,\mathbb{C}}$ of $G_{\mathbb{C}}^{\mathrm{der}}$ is non-trivial. Since $\mathcal{H}$ is compact, K is totally real and at least $2n$ of the factors of $\tilde{G}'_{\mathbb{R}}$ are isomorphic to $\mathrm{SU}_{2,\mathbb{R}}$. Since the corestriction is trivial, it follows that $\tilde{G}'_{\mathbb{R}} \cong \mathrm{SU}_{2,\mathbb{R}}^{2n} \times \mathrm{SL}_{2,\mathbb{R}}$. Since $\mathcal{H}^{\mathrm{der}}$ is compact, it follows that h is conjugate to the map $h_0 : S \longrightarrow G_{\mathbb{R}}$ derived from

$$\tilde{h}_0 : S \longrightarrow \mathrm{GSU}_{2,\mathbb{R}}^{2n} \times \mathrm{GL}_{2,\mathbb{R}} \cong D_{\mathbb{R}}^*, \quad a + b\sqrt{-1} \mapsto \left(1, 1, \cdots, 1, \begin{pmatrix} a & -b \\ b & a \end{pmatrix}\right).$$

This completes the proof. □

6.2 Cyclic lattices with torsion-free quotients

We use this subsection to clarify some of the things from section 4.4. In particular, we discuss the possibilities for the kernel of the map $\mathrm{ev} : X^*(\mathbf{T}) \longrightarrow \overline{\mathbb{Q}}^*$, the image of which is the multiplicative group Φ_{A_v} generated by the eigenvalues of the Frobenius.

Definition 6.1. A sublattice of $\mathbb{Z}^N$ is *cyclic* if it is invariant under the N-cycle $(1\ 2\ 3 \cdots N)$.

We are primarily concerned with cyclic lattices $\mathcal{L} \subseteq \mathbb{Z}^N$ such that the quotient $\mathbb{Z}^N/\mathcal{L}$ is torsion-free. This will help us deduce the possibilities for $\ker(\mathrm{ev})$. We define a map

$$\phi : \mathbb{Z}[X]/(X^N - 1) \longrightarrow \mathbb{Z}^N, \quad \sum_{j=0}^{N-1} a_j X^j \mapsto (a_0, a_1, \cdots, a_{N_1}) \in \mathbb{Z}^N.$$

For an ideal J of $\mathbb{Z}[X]/(X^N - 1)$, the image $\phi(J)$ is a cyclic sublattice of $\mathbb{Z}^N$. Conversely, for a cyclic lattice Λ, $\{\sum_{j=0}^{N-1} a_j X^j : (a_1, \cdots, a_N) \in \Lambda\}$ is an ideal in $\mathbb{Z}[X]/(X^N - 1)$. This gives the bijection

$$\{\text{Cyclic sublattices of } \mathbb{Z}^N\} \longleftrightarrow \{\text{Ideals of } \mathbb{Z}[X]/(X^N - 1)\}.$$

Now, $X^N - 1 = \prod_{d|N} \Phi_d(X)$ where $\Phi_d(X)$ is the d-th cyclotomic polynomial and hence,

$$\mathbb{Z}[X]/(X^N - 1) \cong \prod_{d|N} \mathbb{Z}[\zeta_d]$$

with the integral domains $\mathbb{Z}[\zeta_d]$ integrally closed. So an ideal J of $\mathbb{Z}[X]/(X^N - 1)$ is of the form $\prod_{d|N} J_d$ where J_d is an ideal in $\mathbb{Z}[\zeta_d]$. Furthermore, if the quotient is torsion-free, each ideal J_d is either 0 or the full $\mathbb{Z}[\zeta_d]$. So there are precisely $2^{\tau(N)}$ cyclic sublattices with torsion-free quotient, where $\tau(N)$ is the number of divisors of N.

In particular, if N is a prime, the only ideals of $\mathbb{Z}[X]/(X^N-1) \cong \mathbb{Z} \times \mathbb{Z}[\zeta_N]$ with torsion-free quotients are 0, $\mathbb{Z} \times \mathbb{Z}[\zeta_N]$, $\mathbb{Z} \times 0$ and $0 \times \mathbb{Z}[\zeta_N]$. So the only possibilities for a cyclic sublattice $\mathcal{L} \subseteq \mathbb{Z}^N$ with a torsion-free quotient $\mathbb{Z}^N/\mathcal{L}$ are the following:

(a) 0
(b) $\mathbb{Z}^N$
(c)$\{(x, \cdots, x) : x \in \mathbb{Z}\}$
(d)$\{(x_1, x_2, \cdots, x_N : \sum_{i=1}^{N} x_i = 0\}$.
which verifies the claim in Section 4.

6.3 Central division algebras over totally real fields

We use this subsection to clarify some of the things in Section 5.6. We will need the following lemma.

Lemma 6.2. *Let d be an odd integer. For a finite set of primes $l_1, \cdots, l_r$, there exist infinitely many CM fields E of degree $2d$ over $\mathbb{Q}$ such that:*
(i) *$E/\mathbb{Q}$ is cyclic of degree $2d$*
(ii) *The primes $l_1, \cdots, l_r$ are inert in $E/\mathbb{Q}$.*

Proof. Choose a prime $\mathfrak{p}$ in $\mathbb{Q}(\zeta_d)$ such that:
- $\mathfrak{p}$ is prime to $2d$
- $\mathfrak{p}$ has local degree 1 over $\mathbb{Q}$
- $\mathfrak{p}$ is inert in the degree $2d$ cyclic extension $\mathbb{Q}(\zeta_{4d^2})/\mathbb{Q}(\zeta_d)$
- $\mathfrak{p}$ is split completely in the Kummer extension $\mathbb{Q}\left(\zeta_{2d}, \{\sqrt[2d]{l_1 l_j} : 2 \leq j \leq r\}\right)$ but is inert in the cyclic extension $\mathbb{Q}(\zeta_{2d}, \sqrt[2d]{l_1})$.

Note that by Kummer theory, $\mathbb{Q}(\zeta_{2d}, \sqrt[2d]{l_1}) \cap \mathbb{Q}\left(\zeta_{2d}, \{\sqrt[2d]{l_1 l_j} : 2 \leq j \leq r\}\right) = \mathbb{Q}(\zeta_{2d})$ since $l_1 \notin \langle l_1 l_2, \cdots, l_1 l_r \rangle \subseteq \mathbb{Q}^*$. Hence, the set of primes $\mathfrak{p}$ of $\mathbb{Q}(\zeta_d)$ fulfilling the four conditions has positive density by the Chebotarev density theorem.

By construction, $\mathfrak{p}$ is inert in each of the cyclic extensions $\mathbb{Q}(\zeta_d, \sqrt[2d]{l_j})/\mathbb{Q}(\zeta_d)$ $(1 \leq j \leq r)$. The rational prime p lying under $\mathfrak{p}$ is $\equiv 1 \pmod{2d}$ and $\gcd(\frac{p-1}{2d}, 2d) = 1$. Define E to be the unique degree $2d$ number field contained in $\mathbb{Q}(\zeta_p)$. Since E is of odd index in $\mathbb{Q}(\zeta_p)$, it is a CM field. Each of the rational primes l_i has its inertia degree in $\mathbb{Q}(\zeta_p)$ divisible by $2d$ and since $\gcd(\frac{p-1}{2d}, 2d) = 1$, it follows that every l_i is inert in $E/\mathbb{Q}$. □

We now use this lemma to prove Proposition 4.13 from section 4.6.

Notations: For a field K, $\mathrm{Br}(K)$ is the Brauer group. For an extension L/K, $\mathrm{Br}(L|K)$ is the relative Brauer group, which is the kernel of the restriction map

$$\mathrm{Br}(K) \longrightarrow \mathrm{Br}(L), \quad [D] \mapsto [L \otimes_K D].$$

As before, $\mathcal{S}(L|K)$ is the set of primes of K split completely in L (equivalently, in the Galois closure of L over K).

Proof. (Proposition 4.13) Note that all but finitely many of the fields E constructed in the preceding lemma are linearly disjoint from K. Linear disjointness ensures that $[EK : K] = [E : \mathbb{Q}] = 2d$. So, from Brauer theory, it follows that EK can be embedded in D if and only if EK splits D.

Let $\mathfrak{l}_1, \cdots, \mathfrak{l}_r$ be the non-archimedean primes of K that D is ramified at. Let $l_1, \cdots, l_r$ be the rational primes they lie over. Choose a cyclic extension $E/\mathbb{Q}$ of degree $2d$ inert at the rational primes $l_1, \cdots, l_r$. Since $(2d, [K:\mathbb{Q}]) = 1$, it follows that EK/K is inert at every prime lying over l_i. In particular, it is inert at every $\mathfrak{l}_i$ and hence, $EK_{\mathfrak{l}_i} = K_{\mathfrak{l}_i}{}^{2d}$ for every i. We have the exact sequence

$$0 \longrightarrow \mathrm{Br}(EK|K) \xrightarrow{i} \bigoplus_w \mathrm{Br}(EK^w|K_w) \xrightarrow{\alpha \mapsto \sum_w \mathrm{inv}_w(\alpha)} \frac{1}{2d}\mathbb{Z}/\mathbb{Z} \longrightarrow 0.$$

Now $[D \otimes EK_{\mathfrak{l}_i}] = 0$ in $\mathrm{Br}(EK_{\mathfrak{l}_i})$ for each $\mathfrak{l}_i$. Since EK is a CM field, it follows that $[D] \in \mathrm{Br}(EK|K)$ and $[EK:K] = 2d = \sqrt{[D:K]}$. Hence, EK has an embedding in D. □

We conclude this article with the following Proposition which, when combined with ([No01], Proposition 5.1), helps us construct abelian varieties strongly of Mumford type such that the reduction at some place is simple.

Proposition 6.3. *Let K be a (totally) real field Galois of degree $2n+1$ and let D be a quaternion algebra over K. Then there exists infinitely many CM fields L_1 quadratic over K such that L_1 contains no proper CM subfield and $L_1 \hookrightarrow D$.*

Proof. Let $\mathfrak{l}_1, \cdots, \mathfrak{l}_r$ be the finite primes of K that D is ramified at. Let $K_1, \cdots, K_t$ be the maximal proper subfields of K. For each of them, we choose a prime $\mathfrak{p}_j \in \mathcal{S}(K|K_j)$ co-prime to the primes $\mathfrak{l}_i$ and two distinct primes $\tilde{\mathfrak{p}}_j$, $\tilde{\mathfrak{p}}'_j$ in K lying over $\mathfrak{p}_j$.

Choose a totally positive $\alpha \in K$ such that
- $\mathbb{Q}(\alpha) = K$
- $x^2 + \alpha$ does not split in $(\mathcal{O}_K/\mathfrak{l}_i)[X]$ for any i.
- $x^2 + \alpha$ splits in $(\mathcal{O}_K/\tilde{\mathfrak{p}}_j)[X]$ but is irreducible in $(\mathcal{O}_K/\tilde{\mathfrak{p}}'_j)[X]$ for every j.

The existence of the element α follows from the Chinese remainder theorem. Set $L_1 = K(\sqrt{-\alpha})$. From the third condition, it follows that L_1 has local degree 2 at each $\mathfrak{l}_i$ and since it is totally imaginary, it follows that L_1 splits D and hence, $L_1 \hookrightarrow D$. The third condition implies that L_1 is not Galois over any of the fields K_j, $j = 1, \cdots, t$.

Now suppose L_1 has a proper CM subfield L_0. Then $L_0 \cap K$ is a totally real proper subfield of K of index 2 in L_0. Let $L \cap K \subseteq K_j$ (notations as above). Then $L_1 = L_0 K$ and since $K/\mathbb{Q}$ is Galois, it follows that L_1 is Galois over $L \cap K$ and hence, over K_j. This contradicts the third condition. □

Acknowledgments: The author is grateful to Ben Howard, David Geraghty and Anand Patel for helpful conversations. We also thank Anand Patel, Ananth Shankar and Yuanqing Cai for helpful suggestions on several previous drafts of the article.

References

[Bog80] Bogomolov, *Sur l'algebricite des representations l-adiques*, C. R. Acad. Sci. Paris Ser. A-B 290 (1980), no. 15.

[Chi92] Chi, *l-adic and λ-adic representations associated to abelian varieties defined over number fields*, Amer. J. Math. 114 (1992).

[CCO09] Conrad, Chai, Oort, *CM Liftings*, book manuscript (2011) Chapter 1.

[Del] Deligne, *La Conjecture de Weil I*, Publ. Math. IHES 43 (1974)

[Fal86] Faltings, *Finiteness theorems for abelian varieties over number fields*, Arithmetic geometry (Storrs, Conn., 1984), 1986.

[FS14] Fukshansky and Sun, *On the geometry of cyclic lattices*, Discrete Comput. Geom.,52(2), 2014

[GRR72] Grothendieck, M. Raynaud, and D. S. Rim. *Groupes de monodromie en geometrie algebrique I*, Lecture Notes in Mathematics 288 Springer-Verlag, 1972. Seminaire de Geometrie Algebrique du Bois-Marie 19671969 (SGA 7 I).

[Kne65] Kneser, *Galoiskohomologie halbeinfacher algebraischer Gruppen uber p-adischen Korpern*, Teil II. Math. Zeitschr. 89 (1965).

[Kow03] Kowalski, *Some local-global applications of Kummer theory*, Manuscripta Math., 111(1), 2003.

[Kow06] Kowalski, *Weil numbers generated by other Weil numbers and torsion fields of abelian varieties*, J. London Math. Soc. (2) 74 (2006), no. 2.

[LP95] Larsen and Pink, *Abelian varieties, l-adic representations, and l-independence*, Math. Ann. 302 (1995), no. 3.

[LP97] Larsen and Pink, *A connectedness criterion for l-adic Galois representations*, Israel J. Math. 97 (1997).

[MZ95] Moonen and Zarhin, *Hodge classes and Tate classes on simple abelian fourfolds*, Duke Math. J. 77, 3 (1995).

[Mum69] Mumford, *A note on Shimura's paper "Discontinuous groups and abelian varieties"*, Math. Ann. 181 (1969).

[No95] Noot, *Abelian varieties- Galois representations and properties of ordinary reduction*, Comp. Math. 97 (1995).

[No00] Noot, *Abelian varieties with l-adic Galois representation of Mumford type*, J. Reine Angew. Math. 519 (2000)

[No01] Noot, *On Mumford's families of abelian varieties*, Journal of Pure and Applied Algebra 157 (2001).

[No09] Noot, *Classe de conjugaison du Frobenius dune variete abelienne sur un corps de nombres*, J. Lond. Math. Soc. (2) 79 (2009), no. 1.

[Orr15] Orr, *Lower bounds for ranks of Mumford-Tate groups*, Bull. Soc. Math. France, 143(2), 2015

[Pin98] Pink, *l-adic algebraic monodromy groups, co-characters and the Mumford-Tate conjecture*, J. Reine Angew. Math. 495 (1998

[ST68] Serre and Tate, *Good reduction of abelian varieties*, Ann. of Math. (2), 88, 1968

[Ser79] Serre, *Groupes algebriques associes aux modules de Hodge-Tate*, Journees de Geometrie Algebrique de Rennes. (Rennes, 1978), Vol. III, 1979

[Ser00] Serre, *OEuvres. Collected papers. IV*, Springer-Verlag, Berlin, 2000.

[SZ12] Sheng and Zuo, *On the Newton polygons of abelian varieties of Mumford type*, Preprint

[Tat69] Tate, *Endomorphisms of abelian varieties over finite fields*, Invent. Math, Vol 2, 1966

[Wat69] Waterhouse, *Abelian varieties over finite fields*, Ann. Sci. Ecole Norm. Sup. (4) 2 (1969).

[Win02] Wintenberger, *Demonstration dune conjecture de Lang dans des cas particuliers*, J. Reine Angew. Math. 553 (2002).

[Zh14] Zhao, *On the Mumford-Tate conjecture for abelian fourfolds*, Preprint.

[Zyw14] Zywina, *The splitting of reductions of an abelian variety*, IMRN, vol. 2014, No. 18.

E-mail address: steve.thakur@bc.edu, stevethakur01@gmail.com
Mathematics Department
Boston College.